

Strong on results.



before

Otoscopic view of
tympanic membrane in a patient
who did not respond to ampicillin

after

Same patient after
ten days of Bactrim (trimethoprim
and sulfamethoxazole/Roche) therapy

Simple to take.

in acute otitis media



- ▶ Penetrates and clears middle-ear fluid of susceptible strains of *H. influenzae* and *S. pneumoniae*¹
- ▶ Reduces evidence of inflammation and bulging eardrum²
- ▶ Results in a reduction of fever, pain and other symptoms^{2,3}

Active against 86% of *H. influenzae in vitro*—even amoxicillin- and ampicillin-resistant strains

Overall, 86% of *Haemophilus influenzae* strains taken from sputum cultures prove susceptible *in vitro* to Bactrim.⁴ In one study, 100% of 191 ampicillin-resistant *H. influenzae* isolates were susceptible to Bactrim.⁵ However, *in vitro* data do not necessarily correlate with clinical results.

Active against 91% of *S. pneumoniae in vitro*

In sputum cultures of *Streptococcus pneumoniae*, the most frequent pathogen in acute otitis media, 91% of isolates show susceptibility *in vitro* to Bactrim.⁴

Excellent clinical activity—and economical

In comparative clinical trials in children with acute otitis media, Bactrim *b.i.d.* was unsurpassed by ampicillin, amoxicillin or cefaclor.⁶ And the average cost of Bactrim is lower than that of cefaclor and comparable to that of ampicillin and amoxicillin.⁷ Bactrim is indicated in acute otitis media due to susceptible organisms when it offers an advantage over other antimicrobials. Bactrim is contraindicated in pregnancy, lactation, infants under two months of age and documented megaloblastic anemia due to folate deficiency. To date, there are limited data on the safety of repeated use of Bactrim in children under two years of age.

Cherry-flavored suspension

Bactrim[™] Pediatric

(trimethoprim and sulfamethoxazole/Roche)

B.I.D. for enhanced compliance.



References: 1. Klimek JJ *et al*: *J Pediatr* 96:1087-1089, Jun 1980. 2. Schwartz RH *et al*: *Rev Infect Dis* 4:514-516, Mar-Apr 1982. 3. Cooper J, Inman JS, Dawson AF: *Practitioner* 217:804-809, Nov 1976. 4. Antibiotic Sensitivity Report, Winter 1983. BAC-DATA Medical Information Systems, Inc. 5. Data on file, Hoffmann-La Roche Inc., Nutley, NJ. 6. Wormser GP, Keusch GT, Heel RC: *Drugs* 24:459-518, Dec 1982. 7. *Med Lett Drugs Ther* 23:93-95, Oct 30, 1981.

Please see summary of product information on the following page.

PRESENTS

WILDERNESS MEDICINE

August 12-16, 1985
Lake Tahoe, Nevada

This CME course (23 credit hours in Category I, ACEP, AAFP) has been developed for physicians and other health care professionals who love the outdoors - the open spaces. It will provide the knowledge and skills needed to manage medical problems encountered away from the office, hospital and clinical laboratory.

United Airlines will be offering special airfares, not available to the general public. The meeting will be held at the **Hyatt Lake Tahoe** in Incline Village. A social program has been designed to complement the educational experience.

The fee is \$295.

For brochure and further information, contact:

Office of Continuing Medical Education, M-017B
UCSD School of Medicine
La Jolla, CA 92093
(619)452-3940

*"Come all who need rest and light, bending and breaking with
over work, leave your profits and losses and metallic dividends
and come..."*

John Muir
Dec. 17, 1874

**This Publication
is available in Microform.**



University Microfilms International

Please send additional information

for _____
(name of publication)

Name _____

Institution _____

Street _____

City _____

State _____ Zip _____

300 North Zeeb Road, Dept. P.R., Ann Arbor, Mi. 48106

References: 1. Klimek JJ et al: *J Pediatr* 96:1087-1089, Jun 1980. 2. Schwartz RH et al: *Rev Infect Dis* 4:514-516, Mar-Apr 1982. 3. Cooper J, Inman JS, Dawson AF: *Practitioner* 217:804-809, Nov 1976. 4. Antibiotic Sensitivity Report. Winter 1983. BAC-DATA Medical Information Systems, Inc. 5. Data on file, Hoffmann-La Roche Inc., Nutley, NJ. 6. Wormser GP, Keusch GT, Heel RC: *Drugs* 24:459-518, Dec 1982. 7. *Med Lett Drugs Ther* 23:93-95, Oct 30, 1981.

BACTRIM™ (trimethoprim and sulfamethoxazole/Roche)

Before prescribing, please consult complete product information, a summary of which follows:
Indications and Usage: For the treatment of urinary tract infections due to susceptible strains of the following organisms: *Escherichia coli*, *Klebsiella-Enterobacter*, *Proteus mirabilis*, *Proteus vulgaris*, *Proteus morganii*. It is recommended that initial episodes of uncomplicated urinary tract infections be treated with a single effective antibacterial agent rather than the combination. **Note:** The increasing frequency of resistant organisms limits the usefulness of all antibacterials, especially in these urinary tract infections.

For acute otitis media in children due to susceptible strains of *Haemophilus influenzae* or *Streptococcus pneumoniae* when in physician's judgment it offers an advantage over other antimicrobials. To date, there are limited data on the safety of repeated use of Bactrim in children under two years of age. Bactrim is not indicated for prophylactic or prolonged administration in otitis media at any age. For acute exacerbations of chronic bronchitis in adults due to susceptible strains of *Haemophilus influenzae* or *Streptococcus pneumoniae* when in physician's judgment it offers an advantage over a single antimicrobial agent.

For enteritis due to susceptible strains of *Shigella flexneri* and *Shigella sonnei* when antibacterial therapy is indicated.

Also for the treatment of documented *Pneumocystis carinii* pneumonitis.

Contraindications: Hypersensitivity to trimethoprim or sulfonamides; patients with documented megaloblastic anemia due to folate deficiency; pregnancy at term; nursing mothers because sulfonamides are excreted in human milk and may cause kernicterus; infants less than 2 months of age.

Warnings: BACTRIM SHOULD NOT BE USED TO TREAT STREPTOCOCCAL PHARYNGITIS. Clinical studies show that patients with group A β -hemolytic streptococcal tonsillopharyngitis have higher incidence of bacteriologic failure when treated with Bactrim than do those treated with penicillin. Deaths from hypersensitivity reactions, hepatocellular necrosis, agranulocytosis, aplastic anemia and other blood dyscrasias have been associated with sulfonamides. Experience with trimethoprim is much more limited but occasional interference with hematopoiesis has been reported as well as an increased incidence of thrombopenia with purpura in elderly patients on certain diuretics, primarily thiazides. Sore throat, fever, pallor, purpura or jaundice may be early signs of serious blood disorders. Frequent CBC's are recommended; therapy should be discontinued if a significantly reduced count of any formed blood element is noted.

Precautions: **General:** Use cautiously in patients with impaired renal or hepatic function, possible folate deficiency, severe allergy or bronchial asthma. In patients with glucose-6-phosphate dehydrogenase deficiency, hemolysis, frequently dose-related, may occur. During therapy, maintain adequate fluid intake and perform frequent urinalyses, with careful microscopic examination, and renal function tests, particularly where there is impaired renal function. Bactrim may prolong prothrombin time in those receiving warfarin; reassess coagulation time when administering Bactrim to these patients.

Pregnancy: Teratogenic Effects: Pregnancy Category C. Because trimethoprim and sulfamethoxazole may interfere with folate metabolism, use during pregnancy only if potential benefits justify the potential risk to the fetus.

Adverse Reactions: All major reactions to sulfonamides and trimethoprim are included, even if not reported with Bactrim. **Blood dyscrasias:** Agranulocytosis, aplastic anemia, megaloblastic anemia, thrombopenia, leukopenia, hemolytic anemia, purpura, hypoprothrombinemia and methemoglobinemia. **Allergic reactions:** Erythema multiforme, Stevens-Johnson syndrome, generalized skin eruptions, epidermal necrolysis, urticaria, serum sickness, pruritus, exfoliative dermatitis, anaphylactoid reactions, periorbital edema, conjunctival and scleral injection, photosensitization, arthralgia and allergic myocarditis. **Gastrointestinal reactions:** Glossitis, stomatitis, nausea, emesis, abdominal pains, hepatitis, hepatocellular necrosis, diarrhea, pseudomembranous colitis and pancreatitis. **CNS reactions:** Headache, peripheral neuritis, mental depression, convulsions, ataxia, hallucinations, tinnitus, vertigo, insomnia, apathy, fatigue, muscle weakness and nervousness. **Miscellaneous reactions:** Drug fever, chills, toxic nephrosis with oliguria and anuria, periarteritis nodosa and L.E. phenomenon. Due to certain chemical similarities to some goitrogens, diuretics (acetazolamide, thiazides) and oral hypoglycemic agents, sulfonamides have caused rare instances of goiter production, diuresis and hypoglycemia in patients; cross-sensitivity with these agents may exist. In rats, long-term therapy with sulfonamides has produced thyroid malignancies.

Dosage: Not recommended for infants less than two months of age.

URINARY TRACT INFECTIONS AND SHIGELLOSIS IN ADULTS AND CHILDREN, AND ACUTE OTITIS MEDIA IN CHILDREN:

Adults: Usual adult dosage for urinary tract infections—1 DS tablet (double strength), 2 tablets (single strength) or 4 teasp. (20 ml) b.i.d. for 10-14 days. Use identical daily dosage for 5 days for shigellosis.

Children: Recommended dosage for children with urinary tract infections or acute otitis media—8 mg/kg trimethoprim and 40 mg/kg sulfamethoxazole per 24 hours, in two divided doses for 10 days. Use identical daily dosage for 5 days for shigellosis.

For patients with renal impairment: Use recommended dosage regimen when creatinine clearance is above 30 ml/min. If creatinine clearance is between 15 and 30 ml/min, use one-half the usual regimen. Bactrim is not recommended if creatinine clearance is below 15 ml/min.

ACUTE EXACERBATIONS OF CHRONIC BRONCHITIS IN ADULTS:

Usual adult dosage: 1 DS tablet (double strength), 2 tablets (single strength) or 4 teasp. (20 ml) b.i.d. for 14 days.

PNEUMOCYSTIS CARINII PNEUMONITIS:

Recommended dosage: 20 mg/kg trimethoprim and 100 mg/kg sulfamethoxazole per 24 hours in equal doses every 6 hours for 14 days. See complete product information for suggested children's dosage table.

Supplied: Double Strength (DS) tablets, each containing 160 mg trimethoprim and 800 mg sulfamethoxazole, bottles of 100, 250 and 500; Tel-E-Dose® packages of 100; Prescription Paks of 20. Tablets, each containing 80 mg trimethoprim and 400 mg sulfamethoxazole—bottles of 100 and 500; Tel-E-Dose® packages of 100; Prescription Paks of 40. Pediatric Suspension, containing 40 mg trimethoprim and 200 mg sulfamethoxazole per teaspoonful (5 ml); cherry flavored—bottles of 100 ml and 16 oz (1 pint). Suspension, containing 40 mg trimethoprim and 200 mg sulfamethoxazole per teaspoonful (5 ml); fruit-licorice flavored—bottles of 16 oz (1 pint).



ROCHE LABORATORIES
Division of Hoffmann-La Roche Inc.
Nutley, New Jersey 07110



Team support. Personalized, expert, responsive services, provided every day and when our members need it most.

THE DOCTORS' COMPANY

The Professionals in Professional Liability Insurance


401 WILSHIRE BOULEVARD, SANTA MONICA, CA 90401. (213) 451-3011

**Angina comes in
many forms...**




So does
SORBITRATE®
(ISOSORBIDE DINITRATE)

**Unsurpassed flexibility
in nitrate therapy.**



In hypertension **CORGARD®** (nadolol tablets)



CORGARD® TABLETS Nadolol Tablets

DESCRIPTION: Corgard (nadolol) is a synthetic nonselective beta-adrenergic receptor blocking agent.

CONTRAINDICATIONS: Bronchial asthma, sinus bradycardia and greater than first degree conduction block, cardiogenic shock, and overt cardiac failure (see WARNINGS). **WARNINGS:** **Cardiac Failure**—Sympathetic stimulation may be a vital component supporting circulatory function in congestive heart failure, and its inhibition by beta-blockade may precipitate more severe failure. Although beta-blockers should be avoided in overt congestive heart failure, if necessary, they can be used with caution in patients with a history of failure who are well-compensated, usually with digitalis and diuretics. Beta-adrenergic blocking agents do not abolish the inotropic action of digitalis on heart muscle. **IN PATIENTS WITHOUT A HISTORY OF HEART FAILURE**, continued use of beta-blockers can, in some cases, lead to cardiac failure; therefore, at first sign or symptom of heart failure, digitalize and/or give diuretics, and closely observe response or discontinue nadolol (gradually if possible).

Exacerbation of Ischemic Heart Disease Following Abrupt Withdrawal—Hypersensitivity to catecholamines has been observed in patients withdrawn from beta-blocker therapy; exacerbation of angina and, in some cases, myocardial infarction have occurred after abrupt discontinuation of beta-blocker therapy. In patients with chronic use of nadolol, particularly in patients with chronic angina, gradually reduce dosage over a 1- to 2-week period and carefully monitor the patient. Reinitiate nadolol promptly (at least temporarily) and take other measures appropriate for management of unstable angina if angina markedly worsens or acute coronary insufficiency develops. Warn patients not to interrupt or discontinue therapy without physician's advice. Because coronary artery disease is often asymptomatic and unrecognized, it may be prudent not to discontinue nadolol abruptly in patients treated only for hypertension.

Nonallergic Bronchospasm (e.g., asthma, chronic bronchitis, emphysema)—**PATIENTS WITH BRONCHOSPASTIC DISEASE SHOULD NOT RECEIVE BETA-BLOCKERS.** Administer nadolol with caution since it may inhibit bronchodilation produced by endogenous or exogenous catecholamine stimulation of beta₂ receptors.

Major Surgery—Because beta-blockade impairs the ability of the heart to respond to reflex stimuli and may increase risks of general anesthesia and surgical procedures, resulting in protracted hypotension or low cardiac output, it has generally been suggested that such therapy should be withdrawn several days prior to surgery. Recognition of the increased sensitivity to catecholamines of patients recently withdrawn from beta-blocker therapy, however, has made this recommendation controversial. If possible, withdraw beta-blockers well before surgery takes place. In emergency surgery, inform the anesthesiologist that the patient is on beta-blocker therapy. Use of beta-receptor agonists such as isoproterenol, dopamine, dobutamine, or levaterenol can reverse the effects of nadolol. Difficulty in restarting and maintaining the heart beat has also been reported with beta-adrenergic receptor blocking agents.

Diabetes and Hypoglycemia—Beta-adrenergic blockade may prevent the appearance of premonitory signs and symptoms (e.g., tachycardia and blood pressure changes) of acute hypoglycemia. This is especially important with labile diabetics. Beta-blockade also reduces release of insulin in response to hyperglycemia; therefore, it may be necessary to adjust dose of antidiabetic drugs.

Thyrotoxicosis—Beta-adrenergic blockade may mask certain clinical signs (e.g., tachycardia) of hyperthyroidism. To avoid abrupt withdrawal of beta-adrenergic blockade which might precipitate a thyroid storm, carefully manage patients suspected of developing thyrotoxicosis.

PRECAUTIONS: Impaired Renal Function—Use nadolol with caution (see DOSAGE AND ADMINISTRATION section of package insert).

Information for Patients—Warn patients, especially those with evidence of coronary artery insufficiency, against interruption or discontinuation of nadolol without physician's advice. Although cardiac failure rarely occurs in properly selected patients, advise patients being treated with beta-adrenergic blocking agents to consult physician at first sign of impending failure. Advise patients in event of missed doses.

Drug Interactions—Concurrent administration may result in interactions with: Anesthetics, general—exaggeration of the hypotension induced by general anesthetics (see WARNINGS, Major Surgery). Antidiabetic drugs (oral agents and insulin)—hypoglycemia or hyperglycemia; adjust antidiabetic drug dosage accordingly (see WARNINGS, Diabetes and Hypoglycemia). Catecholamine-depleting drugs (e.g., reserpine)—additive effect; monitor closely for hypotension and/or excessive bradycardia.

Carcinogenesis, Mutagenesis, Impairment of Fertility—In 1 to 2 years' oral toxicologic studies in mice, rats, and dogs, nadolol did not produce significant toxic effects. In 2-year oral carcinogenic studies in rats and mice, nadolol did not produce neoplastic, preneoplastic, or nonneoplastic pathologic lesions.

Pregnancy—In animal reproduction studies with nadolol, evidence of embryo- and fetotoxicity was found in rabbits (but not in rats or hamsters) at doses 5 to 10 times greater (on a mg/kg basis) than maximum indicated human dose; no teratogenic potential was seen in any of these species. There are no well-controlled studies in pregnant women; therefore, use nadolol in pregnant women only if potential benefit justifies potential risk to the fetus. Neonates of mothers who received nadolol at parturition have exhibited bradycardia, hypoglycemia and associated symptoms.

Nursing Mothers—Nadolol is excreted in human milk. Exercise caution when nadolol is administered to a nursing woman.

Pediatric Use—Safety and effectiveness in children have not been established.

ADVERSE REACTIONS: Most adverse effects have been mild and transient and have rarely required nadolol withdrawal.

Cardiovascular—Bradycardia with heart rates of less than 60 beats per minute occurs commonly, and heart rates below 40 beats per minute and/or symptomatic bradycardia were seen in about 2 of 100 patients. Symptoms of peripheral vascular insufficiency, usually of the Raynaud type, have occurred in approximately 2 of 100 patients. Cardiac failure, hypotension, and rhythm/conduction disturbances have each occurred in about 1 of 100 patients. Single instances of first degree and third degree heart block have been reported; intensification of AV block is a known effect of beta-blockers (see also CONTRAINDICATIONS, WARNINGS, and PRECAUTIONS). **Central Nervous System**—Dizziness or fatigue reported in approximately 2 of 100 patients; paresthesias, sedation, and change in behavior reported in approximately 6 of 1000 patients.

Respiratory—Bronchospasm reported in approximately 1 of 1000 patients (see CONTRAINDICATIONS and WARNINGS). **Gastrointestinal**—Nausea, diarrhea, abdominal discomfort, constipation, vomiting, indigestion, anorexia, bloating and flatulence each reported in 1 to 5 of 1000 patients.

Miscellaneous—Each of the following reported in 1 to 5 of 1000 patients: rash; pruritus; headache; dry mouth, eyes, or skin; impotence or decreased libido; facial swelling; weight gain; slurred speech; cough; nasal stuffiness; sweating; tinnitus; blurred vision; infrequent reversible alopecia. The following adverse reactions have been reported in patients taking nadolol and/or other beta-adrenergic blocking agents, but no causal relationship to nadolol has been established. **Central Nervous System**—reversible mental depression progressing to cataplexy; visual disturbances; hallucinations; an acute reversible syndrome characterized by disorientation for time and place; short-term memory loss, emotional lability with slightly clouded sensorium; decreased performance on neuropsychometrics. **Gastrointestinal**—mesenteric arterial thrombosis; ischemic colitis; elevated liver enzymes. **Hematologic**—agranulocytosis; thrombocytopenic or nonthrombocytopenic purpura. **Allergic**—fever combined with rash and sore throat; laryngospasm; respiratory distress. **Miscellaneous**—pernio; urticaria; hypotensive reaction in patients with pheochromocytoma; severe urticaria; Peyronie's disease; the oculomucocutaneous syndrome associated with prolapsed iris has not been reported with nadolol.

Overdosage—Nadolol cannot be removed from the general circulation by hemodialysis. In cases of overdosage, employ the following measures as appropriate. In determining duration of corrective therapy, take into account the duration of effect of nadolol.

Respiratory Distress—Administer a digitalis glycoside and diuretic. It has been reported to vary the dose of digitalis glycoside administered.

Bradycardia—Administer a beta₂-stimulating agent and/or a theophylline derivative.

Cardiac Failure—Administer a beta₂-stimulating agent and/or a theophylline derivative.

DOSAGE: For all patients, DOSAGE MUST BE INDIVIDUALIZED.

For angina pectoris, usual initial dose is 40 mg q.d.; may be gradually increased in 40 to 80 mg increments at 3 to 7 day intervals until optimum clinical response or pronounced slowing of the heart rate; usual maintenance dose is 40 or 80 mg q.d. (doses up to 160 or 240 mg daily may be needed). If treatment is to be discontinued, reduce dosage gradually over a period of 1 to 2 weeks (see WARNINGS).

For hypertension, usual initial dose is 40 mg q.d.; gradually increase in 40 to 80 mg increments until optimum blood pressure reduction is achieved; usual maintenance dose is 40 or 80 mg q.d. (doses up to 240 or 320 mg daily may be needed).

Patients with renal failure require adjustment in dosing interval; see package insert for dosage in these patients.

For full prescribing information, consult package insert.

HOW SUPPLIED: In scored tablets containing 40, 80, 120, or 160 mg nadolol per tablet in bottles of 100 and in Unimatic® unit-dose packs of 100 tablets. The 40 mg, 80 mg, and 120 mg potencies are also available in bottles of 1000 tablets. (J3-527D)

References: 1. Epstein M, Oster JR: Beta-blockers and the kidney. *Min Electrolyte Metab* 8:237-254, 1982. 2. Danesh BJZ, et al: Comparison between short-term renal haemodynamic effects of propranolol and nadolol in essential hypertension: a cross-over study. *Clin Sci* 67:243-248, 1984. 3. Hollenberg NK: Introduction: beta-adrenergic blocking agents—the treatment of hypertension and the kidney. *Royal Soc of Med Int Congress and Symposium Series* 51:1-8, 1982. 4. Frohlich ED, et al: Long-term renal hemodynamic effects of nadolol in patients with essential hypertension. *Am Heart J* 108:1141-1143, 1984. 5. Alexander JC, et al: Long-term experience with nadolol in treatment of hypertension and angina pectoris. *Am Heart J* 108:1136-1140, 1984.



SQUIBB


Innovators in cardiovascular medicine

Since with increasing blood pressure
there may be a progressive decline
in renal blood flow,¹
prescribe . . .

CORGARD[®] (nadolol
tablets)

**Lowers
blood pressure**

**Preserves
renal blood flow**



*Studies prove that CORGARD
(nadolol tablets) preserves renal
blood flow unlike some beta-
blockers, such as propranolol.¹⁻⁴*

*In a two-year study of 106 patients,
CORGARD also decreased serum
creatinine, a measurement of
improved renal function.⁵*

- Offers once-a-day convenience.
- Low incidence of CNS side effects.*
- Avoids potassium depletion.
- Maintains long-term control.


CORGARD[®]
(nadolol tablets)

**STEP-1
FOR HYPERTENSION
WITH ONCE-A-DAY DOSE**

*For a discussion of CONTRAINDICATIONS,
PRECAUTIONS, ADVERSE REACTIONS, and
WARNINGS, including avoidance of abrupt
withdrawal, please see brief summary of prescribing
information on adjacent page.


SQUIBB

Easy To Take



500-mg Pulvules®

250-mg Pulvules

Oral
Suspension
250 mg/5 ml

Oral
Suspension
125 mg/5 ml

Keflex®
cephalexin

Additional information
available to the profession
on request.



420113

Dista Products Company
Division of Eli Lilly and Company
Indianapolis, Indiana 46285
Mfd. by Eli Lilly Industries, Inc.
Carolina, Puerto Rico 00630

PHYSICIANS



We are announcing opportunities for you to serve your country as an Air Force Reserve physician/officer. You can make new professional associations, obtain CME credit and help support the Air Force mission. For those who qualify, retirement credit can be obtained as well as low cost life insurance. One weekend a month plus two weeks a year or less can bring you pride and satisfaction in serving your country.

Call:
(916) 927-0464
(Call Collect)

Or Fill Out Coupon and Mail Today!
To: Health Professions Recruiting
2604 RRS/RSH, McClellan AFB, CA 95652-6002

Name _____
Address _____
City _____ State _____ Zip _____
Phone _____ Prior Service? Yes ____ No ____
Medical Specialty _____ Date of Birth _____

AIR FORCE RESERVE 4-501-1001

A GREAT WAY TO SERVE

1058

Only Ativan[®]

(lorazepam)[®]

effective
rapid
additional
relief of anxiety

relief of anxiety
associated with
depressive symptoms

clearance not
significantly delayed
by age, liver or
kidney dysfunction


cumulative sedative
effects seldom
a problem

low likelihood of
drug interaction
(benzodiazepines produce additive
effects when taken with alcohol or
other CNS depressants)


no significant changes
in vital signs in
cardiovascular patients*

short duration of action,
simple metabolism

*Benzodiazepines have not been shown to
be of benefit in treating the cardiovascular
component.



VASOMOTOR SYMPTOMS THAT DEMAND INTERVENTION




PREMARIN RELIEVES MODERATE TO SEVERE VASOMOTOR SYMPTOMS

Vasomotor symptoms are the most common manifestation of the menopause, affecting up to 75% of menopausal women. Of these, 80% may suffer for more than a year and up to 50% for more than five years. These symptoms can disrupt a woman's life by chronically interrupting sleep, resulting in anxiety and irritability.

In a study of postmenopausal women suffering severe episodes of cutaneous flushing, symptoms improved markedly after administration of estrogen²—the treatment of choice for moderate to severe vasomotor symptoms.³ The estrogen of choice is PREMARIN, the most widely prescribed estrogen for over 40 years. PREMARIN (Conjugated Estrogens Tablets, U.S.P.) relieves moderate to severe vasomotor symptoms of the natural menopause, as well as the acute and often severe symptoms of surgical menopause.

PREMARIN[®]


(CONJUGATED ESTROGENS TABLETS, U.S.P.)



The appearance of these tablets is a trademark of Ayerst Laboratories.

Please see last page for brief summary of full prescribing information.

VAGINAL ATROPHY THAT INTERFERES WITH SEXUALITY




PREMARIN RESTORES THE VAGINAL ENVIRONMENT

In the postmenopausal woman, decreasing levels of estrogen can have devastating effects on a woman's sexual functioning. The pH of vaginal secretions rises, promoting the growth of contaminating organisms. The vaginal epithelium dries and thins, becoming susceptible to irritation, injury, and infection. Sexual relations may be difficult or impossible.



PREMARIN (Conjugated Estrogens, U.S.P.) Vaginal Cream focuses therapy at the site of the problem. Vaginal dryness is relieved, pH reverts to its normal acidity, and the epithelium thickens and becomes more resistant to injury and infection. With the vaginal environment returned to its premenopausal state, sexual function may improve.

PREMARIN[®]

(CONJUGATED ESTROGENS, U.S.P.) Vaginal Cream



POSTMENOPAUSAL BONE LOSS THAT INCAPACITATES




PREMARIN MAY HALT THE DISABLING COURSE OF OSTEOPOROSIS*

Osteoporosis has an enormous epidemiological impact: it affects 10 million American women, and 26% of all women over age 60.³ The disease begins silently and progresses inexorably for 15 to 20 years, until disabling complications occur.⁶

To minimize osteoporotic damage, the condition must be detected early and treated promptly. For many patients, PREMARIN is optimal therapy for osteoporosis, as part of a comprehensive program that includes exercise, good nutrition, and calcium supplements. In a controlled study of postmenopausal and oophorectomized women, PREMARIN (Conjugated Estrogens Tablets, U.S.P.) doses of 0.625 mg/day prevented loss of metacarpal mineral content (see graph above).⁴

PREMARIN[®] (CONJUGATED ESTROGENS TABLETS, U.S.P.)



The appearance of these tablets is a trademark of Ayerst Laboratories.

* Conjugated Estrogens Tablets have been evaluated as probably effective for estrogen-deficiency-induced osteoporosis. Please see last page for brief summary of full prescribing information.

BRIEF SUMMARY (FOR FULL PRESCRIBING INFORMATION AND PATIENT INFORMATION, SEE PACKAGE CIRCULAR)

PREMARIN® Brand of Conjugated Estrogens Tablets, U.S.P.

PREMARIN® Brand of Conjugated Estrogens, U.S.P. Vaginal Cream in a nonliquefying base

1. ESTROGENS HAVE BEEN REPORTED TO INCREASE THE RISK OF ENDOMETRIAL CARCINOMA.

Three independent case control studies have reported an increased risk of endometrial cancer in postmenopausal women exposed to exogenous estrogens for more than one year. This risk was independent of the other known risk factors for endometrial cancer. These studies were further supported by the finding that incidence rates of endometrial cancer have increased sharply since 1969 in eight different areas of the United States with population-based cancer reporting systems, an increase which may be related to the rapidly expanding use of estrogens during the last decade. The three case control studies reported that the risk of endometrial cancer in estrogen users was about 4.5 to 13.9 times greater than in nonusers. The risk appears to depend on both duration of treatment and on estrogen dose. In view of these findings, when estrogens are used for the treatment of menopausal symptoms, the lowest dose that will control symptoms should be utilized and medication should be discontinued as soon as possible. When prolonged treatment is medically indicated, the patient should be reassessed on at least a semiannual basis to determine the need for continued therapy. Although the evidence must be considered preliminary, one study suggests that cyclic administration of low doses of estrogen may carry less risk than continuous administration; it therefore appears prudent to utilize such a regimen. Close clinical surveillance of all women taking estrogens is important. In all cases of undiagnosed persistent or recurring abnormal vaginal bleeding, adequate diagnostic measures should be undertaken to rule out malignancy. There is no evidence at present that "natural" estrogens are more or less hazardous than "synthetic" estrogens at equiestrogenic doses.

2. ESTROGENS SHOULD NOT BE USED DURING PREGNANCY.

The use of female sex hormones, both estrogens and progestogens, during early pregnancy may seriously damage the offspring. It has been shown that females exposed in utero to diethylstilbestrol, a non-steroidal estrogen, have an increased risk of developing in later life a form of vaginal or cervical cancer that is ordinarily extremely rare. This risk has been estimated as not greater than 4 per 1000 exposures. Furthermore, a high percentage of such exposed women (from 30 to 90 percent) have been found to have vaginal adenosis, epithelial changes of the vagina and cervix. Although these changes are histologically benign, it is not known whether they are precursors of malignancy. Although similar data are not available with the use of other estrogens, it cannot be presumed they would not induce similar changes. Several reports suggest an association between intrauterine exposure to female sex hormones and congenital anomalies, including congenital heart defects and limb reduction defects. One case control study estimated a 4.7-fold increased risk of limb reduction defects in infants exposed in utero to sex hormones (estrogens and/or androgens) during the first trimester of pregnancy, or attempted treatment for threatened abortion. Some of these exposures were very short and involved only a few days of treatment. The data suggest that the risk of limb reduction defects in exposed fetuses is somewhat less than 1 per 1000. In the past, female sex hormones have been used during pregnancy in an attempt to treat threatened or habitual abortion. There is considerable evidence that estrogens are ineffective for these indications, and there is no evidence from well controlled studies that progestogens are effective for these uses. If PREMARIN is used during pregnancy or if the patient becomes pregnant while taking this drug, she should be apprised of the potential risks to the fetus, and the advisability of pregnancy continuation.

DESCRIPTION: PREMARIN (Conjugated Estrogens, U.S.P.) contains a mixture of estrogens, obtained exclusively from natural sources, blended to represent the average composition of material derived from pregnant mares' urine. It contains estrone, equilin, and 17 α -dihydroequilin, together with smaller amounts of 17 α -estradiol, equilin, and 17 α -dihydroequilin as salts of their sulfate esters.

INDICATIONS: Based on a review of PREMARIN Tablets by the National Academy of Sciences—National Research Council and/or other information, FDA has classified the indications for use as follows:

Effective: 1. Moderate to severe vasomotor symptoms associated with the menopause. (There is no evidence that estrogens are effective for nervous symptoms or depression without associated vasomotor symptoms, and they should not be used to treat such conditions.)

2. Atrophic vaginitis
3. Kraurosis vulvae
4. Female hypogonadism
5. Female castration
6. Primary ovarian failure
7. Breast cancer (for palliation only) in appropriately selected women and men with metastatic disease.
8. Prostatic carcinoma—palliative therapy of advanced disease.
9. Postpartum breast engorgement—Although estrogens have been widely used for the prevention of postpartum breast engorgement, controlled studies have demonstrated that the incidence of significant painful engorgement in patients not receiving such hormonal therapy is low and usually responsive to appropriate analgesic or other supportive therapy. Consequently, the benefit to be derived from estrogen therapy for this indication must be carefully weighed against the potential increased risk of puerperal thromboembolism associated with the use of large doses of estrogens.

PREMARIN HAS NOT BEEN SHOWN TO BE EFFECTIVE FOR ANY PURPOSE DURING PREGNANCY AND ITS USE MAY CAUSE SEVERE HARM TO THE FETUS (SEE BOXED WARNING).

"Probably" effective: For estrogen deficiency-induced osteoporosis, and only when used in conjunction with other important therapeutic measures such as diet, calcium, physiotherapy, and good general health-promoting measures. Final classification of this indication requires further investigation.

INDICATIONS: PREMARIN (Conjugated Estrogens, U.S.P.) Vaginal Cream is indicated in the treatment of atrophic vaginitis and kraurosis vulvae. PREMARIN Vaginal Cream HAS NOT BEEN SHOWN TO BE EFFECTIVE FOR ANY PURPOSE DURING PREGNANCY AND ITS USE MAY CAUSE SEVERE HARM TO THE FETUS (SEE BOXED WARNING).

CONTRAINDICATIONS: Estrogens should not be used in women (or men) with any of the following conditions: 1. Known or suspected cancer of the breast except in appropriately selected patients being treated for metastatic disease. 2. Known or suspected estrogen-dependent neoplasia. 3. Known or suspected pregnancy (See Boxed Warning). 4. Undiagnosed abnormal genital bleeding. 5. Active thrombophlebitis or thromboembolic disorders. 6. A past history of thrombophlebitis, thrombosis, or thromboembolic disorders associated with previous estrogen use (except when used in treatment of breast or prostatic malignancy).

WARNINGS: Long term continuous administration of natural and synthetic estrogens in certain animal species increases the frequency of carcinomas of the breast, cervix, vagina, and liver. There are now reports that estrogens increase the risk of carcinoma of the endometrium in humans. (See Boxed Warning.) At the present time there is no satisfactory evidence that estrogens given to postmenopausal women increase the risk of cancer of the breast, although a recent study has raised this possibility. There is a need for caution in prescribing estrogens for women with a strong family history of breast cancer or who have breast nodules, fibrocystic disease, or abnormal mammograms. A recent study has reported a 2- to 3-fold increase in the risk of surgically confirmed gallbladder disease in women receiving postmenopausal estrogens.

Adverse effects of oral contraceptives may be expected at the larger doses of estrogen used to treat prostatic or breast cancer or postpartum breast engorgement; it has been shown that there is an increased risk of thrombosis in men receiving estrogens for prostatic cancer and women for postpartum breast engorgement. Users of oral contraceptives have an increased risk of diseases, such as thrombophlebitis, pulmonary embolism, stroke, and myocardial infarction. Cases of retinal thrombosis, mesenteric thrombosis, and optic neuritis have been reported in oral contraceptive users. An increased risk of postsurgery thromboembolic complications has also been reported in users of oral contraceptives. If feasible, estrogen should be discontinued at least 4 weeks before surgery of the type associated with an increased risk of thromboembolism, or during periods of prolonged immobilization. Estrogens should not be used in persons with active thrombophlebitis, thromboembolic disorders, or in persons with a history of such disorders in association with estrogen use. They should be used with caution in patients with cerebral vascular or coronary artery disease. Large doses (5 mg conjugated estrogens per day), comparable to those used to treat cancer of the prostate and breast have been shown to increase the risk of nonfatal myocardial infarction,

pulmonary embolism and thrombophlebitis. When doses of this size are used, any of the thromboembolic and thrombotic adverse effects should be considered a clear risk.

Benign hepatic adenomas should be considered in estrogen users having abdominal pain and tenderness, abdominal mass, or hypovolemic shock. Hepatocellular carcinoma has been reported in women taking estrogen-containing oral contraceptives. Increased blood pressure may occur with use of estrogens in the menopause and blood pressure should be monitored with estrogen use. A worsening of glucose tolerance has been observed in patients on estrogen-containing oral contraceptives. For this reason, diabetic patients should be carefully observed. Estrogens may lead to severe hypercalcemia in patients with breast cancer and bone metastases.

PRECAUTIONS: Physical examination and a complete medical and family history should be taken prior to the initiation of any estrogen therapy with special reference to blood pressure, breasts, abdomen, and pelvic organs, and should include a Papanicolaou smear. As a general rule, estrogen should not be prescribed for longer than one year without another physical examination being performed. Conditions influenced by fluid retention such as asthma, epilepsy, migraine, and cardiac or renal dysfunction, require careful observation. Certain patients may develop manifestations of excessive estrogenic stimulation, such as abnormal or excessive uterine bleeding, mastodynia, etc. Prolonged administration of unopposed estrogen therapy has been reported to increase the risk of endometrial hyperplasia in some patients. Oral contraceptives appear to be associated with an increased incidence of mental depression. Patients with a history of depression should be carefully observed. Preexisting uterine leiomyomata may increase in size during estrogen use. The pathologist should be advised of estrogen therapy when relevant specimens are submitted. If jaundice develops in any patient receiving estrogen, the medication should be discontinued while the cause is investigated. Estrogens should be used with care in patients with impaired liver function, renal insufficiency, metabolic bone diseases associated with hypercalcemia, or in young patients in whom bone growth is not complete.

The following changes may be expected with larger doses of estrogen:

- a. Increased sulfobromophthalein retention.
- b. Increased prothrombin and factors VII, VIII, IX, and X; decreased antithrombin 3; increased norepinephrine-induced platelet aggregability.
- c. Increased thyroid binding globulin (TBG) leading to increased circulating total thyroid hormone, as measured by PBI, T4 by column, or T4 by radioimmunoassay. Free T3 resin uptake is decreased, reflecting the elevated TBG; free T4 concentration is unaltered.
- d. Impaired glucose tolerance.
- e. Decreased pregnandiol excretion.
- f. Reduced response to metyrapone test.
- g. Reduced serum folate concentration.
- h. Increased serum triglyceride and phospholipid concentration.

As a general principle, the administration of any drug to nursing mothers should be done only when clearly necessary since many drugs are excreted in human milk.

ADVERSE REACTIONS: The following have been reported with estrogenic therapy, including oral contraceptives: breakthrough bleeding, spotting, change in menstrual flow, dysmenorrhea; premenstrual-like syndrome; amenorrhea during and after treatment; increase in size of uterine fibromyoma; vaginal candidiasis; change in cervical erosion and in degree of cervical secretion; cystitis-like syndrome; tenderness, enlargement, secretion (of breasts); nausea, vomiting, abdominal cramps, bloating; cholestatic jaundice; chloasma or melasma which may persist when drug is discontinued; erythema multiforme; erythema nodosum; hemorrhagic eruption; loss of scalp hair; hirsutism; steepening of corneal curvature; intolerance to contact lenses; headache, migraine, dizziness, mental depression, chorea; increase or decrease in weight; reduced carbohydrate tolerance; aggravation of porphyria; edema; changes in libido.

ACUTE OVERDOSAGE: May cause nausea, and withdrawal bleeding may occur in females.

DOSEAGE AND ADMINISTRATION:

PREMARIN® Brand of Conjugated Estrogens Tablets, U.S.P.

1. Given cyclically for short-term use only. For treatment of moderate to severe vasomotor symptoms, atrophic vaginitis, or kraurosis vulvae associated with the menopause (0.3 to 1.25 mg or more daily).

The lowest dose that will control symptoms should be chosen and medication should be discontinued as promptly as possible.

Administration should be cyclic (e.g., three weeks on and one week off).

Attempts to discontinue or taper medication should be made at three to six month intervals.

2. Given cyclically: Female hypogonadism. Female castration. Primary ovarian failure. Osteoporosis.

Female hypogonadism—2.5 to 7.5 mg daily, in divided doses for 20 days, followed by a rest period of 10 days; duration. If bleeding does not occur by the end of this period, the same dosage schedule is repeated. The number of courses of estrogen therapy necessary to produce bleeding may vary depending on the responsiveness of the endometrium.

If bleeding occurs before the end of the 10 day period, begin a 20 day estrogen-progestin cyclic regimen with PREMARIN (Conjugated Estrogens Tablets, U.S.P.), 2.5 to 7.5 mg daily in divided doses, for 20 days. During the last five days of estrogen therapy, give an oral progestin. If bleeding occurs before this regimen is concluded, therapy is discontinued and may be resumed on the fifth day of bleeding.

Female castration and primary ovarian failure—1.25 mg daily, cyclically. Adjust upward or downward according to response of the patient. For maintenance, adjust dosage to lowest level that will provide effective control.

Osteoporosis (to retard progression)—1.25 mg daily, cyclically.

3. Given for a few days: Prevention of postpartum breast engorgement—3.75 mg every four hours for five days, or 1.25 mg every four hours for five days.

4. Given chronically: Inoperable progressing prostatic cancer—1.25 to 2.5 mg three times daily. Inoperable progressing breast cancer in appropriately selected men and postmenopausal women—10 mg three times daily for a period of at least three months.

Patients with an intact uterus should be monitored for signs of endometrial cancer and appropriate measures taken to rule out malignancy in the event of persistent or recurring abnormal vaginal bleeding.

PREMARIN® Brand of Conjugated Estrogens, U.S.P. Vaginal Cream

Given cyclically for short-term use only. For treatment of atrophic vaginitis or kraurosis vulvae.

The lowest dose that will control symptoms should be chosen and medication should be discontinued as promptly as possible.

Administration should be cyclic (e.g., three weeks on and one week off).

Attempts to discontinue or taper medication should be made at three to six month intervals.

Usual dosage range: 2 to 4 g daily, intravaginally or topically, depending on the severity of the condition.

Treated patients with an intact uterus should be monitored closely for signs of endometrial cancer and appropriate diagnostic measures should be taken to rule out malignancy in the event of persistent or recurring abnormal vaginal bleeding.

HOW SUPPLIED: PREMARIN (Conjugated Estrogens Tablets, U.S.P.). No. 865—Each purple tablet contains 2.5 mg in bottles of 100 and 1,000. No. 866—Each yellow tablet contains 1.25 mg in bottles of 100 and 1,000. Also in Cycle Pack of 21 and in unit dose package of 100. No. 864—Each white tablet contains 0.9 mg in bottles of 100. Also in Cycle Pack of 21. No. 867—Each maroon tablet contains 0.625 mg in bottles of 100 and 1,000. Also in Cycle Pack of 21 and unit dose package of 100. No. 868—Each green tablet contains 0.3 mg in bottles of 100 and 1,000. The appearance of these tablets is a trademark of Ayerst Laboratories.

PREMARIN (Conjugated Estrogens, U.S.P.) Vaginal Cream—No. 872—Each gram contains 0.625 mg Conjugated Estrogens, U.S.P. (Also contains cetyl esters wax, cetyl alcohol, white wax, glyceryl monostearate, propylene glycol monostearate, methyl stearate, phenylethyl alcohol, sodium lauryl sulfate, glycerin, and mineral oil.)

Combination package: Each contains Net Wt. 1½ oz. (42.5 g) tube with one calibrated plastic applicator.

Also Available—Refill package: Each contains Net Wt. 1½ oz. (42.5 g) tube.

4340R2/785

REFERENCES: 1. Judd HL: After the menopause. *Transition* 1983;1:19-30. 2. Ertik Y, Tatarin IV, Meldrum DR, et al: Association of waking episodes with menopausal hot flushes. *JAMA* 1981;245:1741-1744. 3. Meldrum DR: The pathophysiology of postmenopausal symptoms. *Sem Reprod Endocrinol* 1983;1(Feb):11-17. 4. Lindsay R, Hart DM, Clark DM: The minimum effective dose of estrogen for prevention of postmenopausal bone loss. *Obstet Gynecol* 1984;63:759-763. 5. Katz WA: *Rheumatic Diseases: Diagnosis and Management*. Philadelphia, JB Lippincott Co, 1977, p 672. 6. Reese WD: A better way to screen for osteoporosis. *Contemp Ob/Gyn* 1983;22(November):116-131.

Ayerst. AYERST LABORATORIES
New York, N.Y. 10017

© 1985 Ayerst Laboratories.

SCPIE.

An enviable record of healthy growth and achievement.

In Perfect Health!

That's the professional opinion on SCPIE.

By every key test, SCPIE is strong...

Best's, the highly-respected, independent, national insurance rating bureau, has just given SCPIE its first rating, "A" (Excellent).

Assets: SCPIE has more assets than any other physician-owned company in California.

Reserves: SCPIE has reserves to pay all outstanding and anticipated claims.

Surplus: SCPIE maintains an adequate surplus as a cushion against unexpected losses.

Membership: SCPIE is strong and growing, with more members in Southern California than any other company.

Management: SCPIE's all-physician Board of Governors and professional insurance staff have proven that conservative management pays off.

Underwriting: Physician control helps keep losses down.

Why is SCPIE the leader?

SCPIE has the lowest overhead of any physician-owned company in Southern California.

For the sixth year, SCPIE is repaying the Capital Contributions which physicians put up to start the company.

Returns of Capital Contributions are approved by the State Department of Insurance—further confirmation of SCPIE's strength.

SCPIE returns premium and investment income not needed to pay claims and expenses. In 1985 SCPIE is returning \$7 million as Experience Credits for policy years 1976-81. This makes significant reductions in net premiums.

SCPIE is here to stay.

SCPIE has become the seventh largest writer of medical liability coverage in the entire nation. An enviable record of healthy growth and achievement.


scpie

Southern California
Physicians Insurance
Exchange

2029 Century Park East
Suite 2300
Los Angeles, CA 90067
(213) 552-8900



Right from the start
in hypertension...



Once-daily **INDERAL LA** (propranolol HCl) for smooth blood pressure control without the potassium problems of diuretics

Once-daily **INDERAL LA** (propranolol HCl) avoids the risk of diuretic-induced ECG abnormalities due to hypokalemia.^{1,2} In addition, **INDERAL LA** preserves potassium balance without additive agents or supplements while providing simple, well-tolerated therapy with broad cardiovascular benefits.

Once-daily **INDERAL LA** for the cardiovascular benefits of the world's leading beta blocker

Simply start with 80 mg once daily. Dosage may be increased to 120 mg to 160 mg once daily as needed to achieve additional control.

Like conventional **INDERAL** tablets, **INDERAL LA** should not be used in the presence of congestive heart failure, sinus bradycardia, heart block greater than first degree, and bronchial asthma.




80 mg 120 mg 160 mg

The appearance of these capsules is a registered trademark of Ayerst Laboratories.

Please see brief summary of prescribing information on the next page for further details.

Once-daily For beta-1/beta-2 blockade **INDERAL[®] LA** (PROPRANOLOL HCl) LONG ACTING CAPSULES



The appearance of these capsules is a registered trademark of Ayerst Laboratories.

BRIEF SUMMARY (FOR FULL PRESCRIBING INFORMATION, SEE PACKAGE CIRCULAR.)

INDERAL[®] LA brand of propranolol hydrochloride (Long Acting Capsules)
DESCRIPTION. Inderal LA is formulated to provide a sustained release of propranolol hydrochloride. Inderal LA is available as 80 mg, 120 mg, and 160 mg capsules.

CLINICAL PHARMACOLOGY. Inderal is a nonselective beta-adrenergic receptor blocking agent possessing no other autonomic nervous system activity. It specifically competes with beta-adrenergic receptor stimulating agents for available receptor sites. When access to beta-receptor sites is blocked by Inderal, the chronotropic, inotropic, and vasodilator responses to beta-adrenergic stimulation are decreased proportionately.

INDERAL LA Capsules (80, 120, and 160 mg) release propranolol HCl at a controlled and predictable rate. Peak blood levels following dosing with Inderal LA occur at about 6 hours and the apparent plasma half-life is about 10 hours. When measured at steady state over a 24-hour period the areas under the propranolol plasma concentration-time curve (AUCs) for the capsules are approximately 60% to 65% of the AUCs for a comparable divided daily dose of Inderal tablets. The lower AUCs for the capsules are due to greater hepatic metabolism of propranolol, resulting from the slower rate of absorption of propranolol. Over a twenty-four (24) hour period, blood levels are fairly constant for about twelve (12) hours then decline exponentially.

INDERAL LA should not be considered a simple mg for mg substitute for conventional propranolol and the blood levels achieved do not match (are lower than) those of two to four times daily dosing with the same dose. When changing to Inderal LA from conventional propranolol, a possible need for retitration upwards should be considered especially to maintain effectiveness at the end of the dosing interval. In most clinical settings, however, such as hypertension or angina where there is little correlation between plasma levels and clinical effect, Inderal LA has been therapeutically equivalent to the same mg dose of conventional Inderal as assessed by 24-hour effects on blood pressure and on 24-hour exercise responses of heart rate, systolic pressure and rate pressure product. Inderal LA can provide effective beta blockade for a 24-hour period.

The mechanism of the antihypertensive effect of Inderal has not been established. Among the factors that may be involved in contributing to the antihypertensive action are (1) decreased cardiac output, (2) inhibition of renin release by the kidneys, and (3) diminution of tonic sympathetic nerve outflow from vasomotor centers in the brain. Although total peripheral resistance may increase initially, it readjusts to or below the pretreatment level with chronic use. Effects on plasma volume appear to be minor and somewhat variable. Inderal has been shown to cause a small increase in serum potassium concentration when used in the treatment of hypertensive patients.

In angina pectoris, propranolol generally reduces the oxygen requirement of the heart at any given level of effort by blocking the catecholamine-induced increases in the heart rate, systolic blood pressure, and the velocity and extent of myocardial contraction. Propranolol may increase oxygen requirements by increasing left ventricular fiber length, end diastolic pressure and systolic ejection period. The net physiologic effect of beta-adrenergic blockade is usually advantageous and is manifested during exercise by delayed onset of pain and increased work capacity.

In dosages greater than required for beta blockade, Inderal also exerts a quinidine-like or anesthetic-like membrane action which affects the cardiac action potential. The significance of the membrane action in the treatment of arrhythmias is uncertain.

The mechanism of the antimigraine effect of propranolol has not been established. Beta-adrenergic receptors have been demonstrated in the pial vessels of the brain.

Beta receptor blockade can be useful in conditions in which, because of pathologic or functional changes, sympathetic activity is detrimental to the patient. But there are also situations in which sympathetic stimulation is vital. For example, in patients with severely damaged hearts, adequate ventricular function is maintained by virtue of sympathetic drive which should be preserved. In the presence of AV block, greater than first degree, beta blockade may prevent the necessary facilitating effect of sympathetic activity on conduction. Beta blockade results in bronchial constriction by interfering with adrenergic bronchodilator activity which should be preserved in patients subject to bronchospasm.

Propranolol is not significantly dialyzable.

INDICATIONS AND USAGE. **Hypertension:** Inderal LA is indicated in the management of hypertension; it may be used alone or used in combination with other antihypertensive agents, particularly a thiazide diuretic. Inderal LA is not indicated in the management of hypertensive emergencies.

Angina Pectoris Due to Coronary Atherosclerosis: Inderal LA is indicated for the long-term management of patients with angina pectoris.

Migraine: Inderal LA is indicated for the prophylaxis of common migraine headache. The efficacy of propranolol in the treatment of a migraine attack that has started has not been established and propranolol is not indicated for such use.

Hypertrophic Subaortic Stenosis: Inderal LA is useful in the management of hypertrophic subaortic stenosis, especially for treatment of exertional or other stress-induced angina, palpitations, and syncope. Inderal LA also improves exercise performance. The effectiveness of propranolol hydrochloride in this disease appears to be due to a reduction of the elevated outflow pressure gradient which is exacerbated by beta-receptor stimulation. Clinical improvement may be temporary.

CONTRAINDICATIONS. Inderal is contraindicated in 1) cardiogenic shock, 2) sinus bradycardia and greater than first degree block, 3) bronchial asthma, 4) congestive heart failure (see WARNINGS) unless the failure is secondary to a tachyarrhythmia treatable with Inderal.

WARNINGS. **CARDIAC FAILURE:** Sympathetic stimulation may be a vital component supporting circulatory function in patients with congestive heart failure, and its inhibition by beta blockade may precipitate more severe failure. Although beta blockers should be avoided in overt congestive heart failure, if necessary, they can be used with close follow-up in patients with a history of failure who are well compensated and are receiving digitalis and diuretics. Beta-adrenergic blocking agents do not abolish the inotropic action of digitalis on heart muscle.

IN PATIENTS WITHOUT A HISTORY OF HEART FAILURE, continued use of beta blockers can, in some cases, lead to cardiac failure. Therefore, at the first sign or symptom of heart failure, the patient should be digitalized and/or treated with diuretics, and the response observed closely, or Inderal should be discontinued (gradually, if possible).

IN PATIENTS WITH ANGINA PECTORIS, there have been reports of exacerbation of angina and, in some cases, myocardial infarction, following abrupt discontinuance of Inderal therapy. Therefore, when discontinuance of Inderal is planned the dosage should be gradually reduced over at least a few weeks, and the patient should be cautioned against interruption or cessation of therapy without the physician's advice. If Inderal therapy is interrupted and exacerbation of angina occurs, it is usually advisable to reinstitute Inderal therapy and take other measures appropriate for the management of unstable angina pectoris. Since coronary artery disease may be unrecognized, it may be prudent to follow the above advice in patients considered at risk of having occult atherosclerotic heart disease who are given propranolol for other indications.

Nonallergic Bronchospasm (e.g., chronic bronchitis, emphysema)—PATIENTS WITH BRONCHOSPASTIC DISEASES SHOULD IN GENERAL NOT RECEIVE BETA BLOCKERS. Inderal should be administered with caution since it may block bronchodilation produced by endogenous and exogenous catecholamine stimulation of beta receptors.

MAJOR SURGERY: The necessity or desirability of withdrawal of beta-blocking therapy prior to major surgery is controversial. It should be noted, however, that the impaired ability of the heart to respond to reflex adrenergic stimuli may augment the risks of general anesthesia and surgical procedures.

INDERAL (propranolol HCl), like other beta blockers, is a competitive inhibitor of beta-receptor agonists and its effects can be reversed by administration of such agents, e.g., dobutamine or isoproterenol. However, such patients may be subject to protracted severe hypotension. Difficulty in starting and maintaining the heartbeat has also been reported with beta blockers.

DIABETES AND HYPOGLYCEMIA: Beta-adrenergic blockade may prevent the appearance of certain premonitory signs and symptoms (pulse rate and pressure changes) of acute hypoglycemia in labile insulin-dependent diabetes. In these patients, it may be more difficult to adjust the dosage of insulin.

THYROTOXICOSIS: Beta blockade may mask certain clinical signs of hyperthyroidism. Therefore, abrupt withdrawal of propranolol may be followed by an exacerbation of symptoms of hyperthyroidism, including thyroid storm. Propranolol does not distort thyroid function tests.

IN PATIENTS WITH WOLFF-PARKINSON-WHITE SYNDROME, several cases have been reported in which, after propranolol, the tachycardia was replaced by a severe bradycardia requiring a demand pacemaker. In one case this resulted after an initial dose of 5 mg propranolol.

PRECAUTIONS. **General:** Propranolol should be used with caution in patients with impaired hepatic or renal function. Inderal (propranolol HCl) is not indicated for the treatment of hypertensive emergencies.

Beta adrenoceptor blockade can cause reduction of intraocular pressure. Patients should be told that Inderal may interfere with the glaucoma screening test. Withdrawal may lead to a return of increased intraocular pressure.

Clinical Laboratory Tests: Elevated blood urea levels in patients with severe heart disease, elevated serum transaminase, alkaline phosphatase, lactate dehydrogenase.

DRUG INTERACTIONS: Patients receiving catecholamine-depleting drugs such as reserpine should be closely observed if Inderal is administered. The added catecholamine-blocking action may produce an excessive reduction of resting sympathetic nervous activity which may result in hypotension, marked bradycardia, vertigo, syncopal attacks, or orthostatic hypotension.

Carcinogenesis, Mutagenesis, Impairment of Fertility: Long-term studies in animals have been conducted to evaluate toxic effects and carcinogenic potential. In 18-month studies in both rats and mice, employing doses up to 150 mg/kg/day, there was no evidence of significant drug-induced toxicity. There were no drug-related tumorigenic effects at any of the dosage levels. Reproductive studies in animals did not show any impairment of fertility that was attributable to the drug.

Pregnancy: Pregnancy Category C. Inderal has been shown to be embryotoxic in animal studies at doses about 10 times greater than the maximum recommended human dose.

There are no adequate and well-controlled studies in pregnant women. Inderal should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers: Inderal is excreted in human milk. Caution should be exercised when Inderal is administered to a nursing woman.

Pediatric Use: Safety and effectiveness in children have not been established.

ADVERSE REACTIONS. Most adverse effects have been mild and transient and have rarely required the withdrawal of therapy.

Cardiovascular: bradycardia; congestive heart failure; intensification of AV block; hypotension; paresthesia of hands; thrombocytopenic purpura; arterial insufficiency, usually of the Raynaud type.

Central Nervous System: lightheadedness; mental depression manifested by insomnia, lassitude, weakness, fatigue; reversible mental depression progressing to cataplexy; visual disturbances, hallucinations; an acute reversible syndrome characterized by disorientation for time and place, short-term memory loss, emotional lability, slightly clouded sensorium, and decreased performance on neuropsychometric tests.

Gastrointestinal: nausea, vomiting, epigastric distress, abdominal cramping, diarrhea, constipation, mesenteric arterial thrombosis, ischemic colitis.

Allergic: pharyngitis and agranulocytosis, erythematous rash, fever combined with aching and sore throat, laryngospasm and respiratory distress.

Respiratory: bronchospasm.

Hematologic: agranulocytosis, nonthrombocytopenic purpura, thrombocytopenic purpura.

Auto-Immune: In extremely rare instances, systemic lupus erythematosus has been reported.

Miscellaneous: alopecia, LE-like reactions, psoriasisiform rashes, dry eyes, male impotence, and Peyronie's disease have been reported rarely. Oculomucocutaneous reactions involving the skin, serous membranes and conjunctivae reported for a beta blocker (practolol) have not been associated with propranolol.

DOSEAGE AND ADMINISTRATION. Inderal LA provides propranolol hydrochloride in a sustained-release capsule for administration once daily. If patients are switched from Inderal tablets to Inderal LA capsules, care should be taken to assure that the desired therapeutic effect is maintained. Inderal LA should not be considered a simple mg for mg substitute for Inderal. Inderal LA has different kinetics and produces lower blood levels. Retitration may be necessary especially to maintain effectiveness at the end of the 24-hour dosing interval.

HYPERTENSION—Dosage must be individualized. The usual initial dosage is 80 mg Inderal LA once daily, whether used alone or added to a diuretic. The dosage may be increased to 120 mg once daily or higher until adequate blood pressure control is achieved. The usual maintenance dosage is 120 to 160 mg once daily. In some instances a dosage of 640 mg may be required. The time needed for full hypertensive response to a given dosage is variable and may range from a few days to several weeks.

ANGINA PECTORIS—Dosage must be individualized. Starting with 80 mg Inderal LA once daily, dosage should be gradually increased at three to seven day intervals until optimum response is obtained. Although individual patients may respond at any dosage level, the average optimum dosage appears to be 160 mg once daily. In angina pectoris, the value and safety of dosage exceeding 320 mg per day have not been established.

If treatment is to be discontinued, reduce dosage gradually over a period of a few weeks (see WARNINGS).

MIGRAINE—Dosage must be individualized. The initial oral dose is 80 mg Inderal LA once daily. The usual effective dose range is 160-240 mg once daily. The dosage may be increased gradually to achieve optimum migraine prophylaxis. If a satisfactory response is not obtained within four to six weeks after reaching the maximum dose, Inderal LA therapy should be discontinued. It may be advisable to withdraw the drug gradually over a period of several weeks.

HYPERTROPHIC SUBAORTIC STENOSIS—80-160 mg Inderal LA once daily.
PEDIATRIC DOSAGE—At this time the data on the use of the drug in this age group are too limited to permit adequate directions for use.


REFERENCES

- Holland OB, Nixon JV, Kuhnert L: Diuretic-induced ventricular ectopic activity. *Am J Med* 1981;70:762-768.
- Holme I, Helgeland A, Hjermann I, et al: Treatment of mild hypertension with diuretics. The importance of ECG abnormalities in the Oslo study and in MRFIT. *JAMA* 1984;251:1298-1299.

Ayerst[®] AYERST LABORATORIES
New York, N.Y. 10017

9411/1184

New
Motrin[®] 800 mg
ibuprofen



a stronger reason
to prescribe
Motrin Tablets

New **Motrin® 800** TABLETS mg ibuprofen

Extra-Strength Motrin Tablets— a convenient way to tap the full potential of Motrin:

The newest strength of *Motrin* Tablets

makes treatment easier for arthritis patients who need the doses that provide higher levels of anti-inflammatory activity as well as potent analgesia...just 1 tablet t.i.d. provides 2400 mg/day.

expands the dosage convenience of MOTRIN Tablets—makes it even easier to adjust the dosage of MOTRIN to each patient's needs...the new dosage range of up to 3200 mg/day can be achieved on a q.i.d. regimen. Gastroscopic studies at varying doses show an increased tendency toward gastric irritation at higher doses. However, at comparable doses, gastric irritation is about half that seen with aspirin.

provides economy...patients should pay less for MOTRIN Tablets than comparable dosages of Clinoril, Feldene, or Naprosyn.

provides, above all, the experience-proven efficacy and safety profile of *Motrin*. MOTRIN continues to be America's most often prescribed nonsteroidal anti-inflammatory agent.

Please turn the page for a brief summary of prescribing information.

Upjohn

The Upjohn Company, Kalamazoo, Michigan 49001



“When it comes to cardiovascular medicine, I like to know exactly what my patients are swallowing.”

There are doctors who say that generic drugs have a place in their practice—but not necessarily in the treatment of serious or potentially life-threatening disease. And when they consider that the average patient pays only about 45¢ a day for INDERAL (propranolol HCl) Tablets, there's not much left to discuss.

When it's INDERAL Tablets you want for the treatment of hypertension, angina, arrhythmias, or post-MI patients, make sure you specify “Dispense As Written” (DAW), “Do Not Substitute,” or whatever is required in your State. That way, you'll know exactly what your patients will get.


Please see next page for brief summary of prescribing information.



"When it comes to cardiovascular medicine, I like to know exactly what my patients are swallowing."

INDERAL® Tablets

BRAND OF PROPRANOLOL HCl



BRIEF SUMMARY (FOR FULL PRESCRIBING INFORMATION, SEE PACKAGE CIRCULAR.)

INDERAL® (propranolol hydrochloride) Tablets

CONTRAINDICATIONS

INDERAL is contraindicated in 1) cardiogenic shock, 2) sinus bradycardia and greater than first degree block, 3) bronchial asthma, 4) congestive heart failure (see WARNINGS) unless the failure is secondary to a tachyarrhythmia treatable with INDERAL.

WARNINGS

CARDIAC FAILURE: Sympathetic stimulation may be a vital component supporting circulatory function in patients with congestive heart failure, and its inhibition by beta blockade may precipitate more severe failure. Although beta blockers should be avoided in overt congestive heart failure, if necessary they can be used with close follow-up in patients with a history of failure who are well compensated and are receiving digitalis and diuretics. Beta-adrenergic blocking agents do not abolish the inotropic action of digitalis on heart muscle.

IN PATIENTS WITHOUT A HISTORY OF HEART FAILURE, continued use of beta blockers can, in some cases, lead to cardiac failure. Therefore, at the first sign or symptom of heart failure, the patient should be digitalized and/or treated with diuretics, and the response observed closely, or INDERAL should be discontinued (gradually, if possible).

IN PATIENTS WITH ANGINA PECTORIS, there have been reports of exacerbation of angina and, in some cases, myocardial infarction, following abrupt discontinuance of INDERAL therapy. Therefore, when discontinuance of INDERAL is planned the dosage should be gradually reduced over at least a few weeks and the patient should be cautioned against interruption or cessation of therapy without the physician's advice. If INDERAL therapy is interrupted and exacerbation of angina occurs, it usually is advisable to reinstitute INDERAL therapy and take other measures appropriate for the management of unstable angina pectoris. Since coronary artery disease may be unrecognized, it may be prudent to follow the above advice in patients considered at risk of having occult atherosclerotic heart disease who are given propranolol for other indications.

Nonallergic Bronchospasm (e.g., chronic bronchitis, emphysema)—PATIENTS WITH BRONCHOSPASTIC DISEASES SHOULD IN GENERAL NOT RECEIVE BETA BLOCKERS. INDERAL should be administered with caution since it may block bronchodilation produced by endogenous and exogenous catecholamine stimulation of beta receptors.

MAJOR SURGERY: The necessity or desirability of withdrawal of beta-blocking therapy prior to major surgery is controversial. It should be noted, however, that the impaired ability of the heart to respond to reflex adrenergic stimuli may augment the risks of general anesthesia and surgical procedures.

INDERAL, like other beta blockers, is a competitive inhibitor of beta-receptor agonists and its effects can be reversed by administration of such agents, e.g., dobutamine or isoproterenol. However, such patients may be subject to protracted severe hypotension. Difficulty in starting and maintaining the heartbeat has also been reported with beta blockers.

DIABETES AND HYPOGLYCEMIA: Beta-adrenergic blockade may prevent the appearance of certain premonitory signs and symptoms (pulse rate and pressure changes) of acute hypoglycemia in labile insulin-dependent diabetes. In these patients, it may be more difficult to adjust the dosage of insulin.

THYROTOXICOSIS: Beta blockade may mask certain clinical signs of hyperthyroidism. Therefore, abrupt withdrawal of propranolol may be followed by an exacerbation of symptoms of hyperthyroidism, including thyroid storm. Propranolol does not distort thyroid function tests.

IN PATIENTS WITH WOLFF-PARKINSON-WHITE SYNDROME, several cases have been reported in which, after propranolol, the tachycardia was replaced by a severe bradycardia requiring a demand pacemaker. In one case this resulted after an initial dose of 5 mg propranolol.

PRECAUTIONS

General: Propranolol should be used with caution in patients with impaired hepatic or renal function. INDERAL is not indicated for the treatment of hypertensive emergencies.

Beta-adrenoreceptor blockade can cause reduction of intraocular pressure. Patients should be told that INDERAL (propranolol hydrochloride) may interfere with the glaucoma screening test. Withdrawal may lead to a return of increased intraocular pressure.

Clinical Laboratory Tests: Elevated blood urea levels in patients with severe heart disease, elevated serum transaminase, alkaline phosphatase, lactate dehydrogenase.

DRUG INTERACTIONS: Patients receiving catecholamine-depleting drugs such as reserpine should be closely observed if INDERAL is administered. The added catecholamine-blocking action may produce an excessive reduction of resting sympathetic nervous activity which may result in hypotension, marked bradycardia, vertigo, syncopal attacks, or orthostatic hypotension.

Carcinogenesis, Mutagenesis, Impairment of Fertility: Long-term studies in animals have been conducted to evaluate toxic effects and carcinogenic potential. In 18-month studies in both rats and mice, employing doses up to 150 mg/kg/day, there was no evidence of significant drug-induced toxicity. There were no drug-related tumorigenic effects at any of the dosage levels. Reproductive studies in animals did not show any impairment of fertility that was attributable to the drug.

Pregnancy: Pregnancy Category C. INDERAL has been shown to be embryotoxic in animal studies at doses about 10 times greater than the maximum recommended human dose.

There are no adequate and well-controlled studies in pregnant women. INDERAL should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers: INDERAL is excreted in human milk. Caution should be exercised when INDERAL is administered to a nursing woman.

Pediatric Use: Safety and effectiveness in children have not been established.

ADVERSE REACTIONS

Most adverse effects have been mild and transient and have rarely required the withdrawal of therapy.

Cardiovascular: bradycardia; congestive heart failure; intensification of AV block; hypotension; paresthesia of hands; thrombocytopenic purpura; arterial insufficiency, usually of the Raynaud type.

Central Nervous System: Lightheadedness; mental depression manifested by insomnia, lassitude, weakness, fatigue; reversible mental depression progressing to catatonia; visual disturbances; hallucinations; an acute reversible syndrome characterized by disorientation for time and place, short-term memory loss, emotional lability, slightly clouded sensorium, and decreased performance on neuropsychometrics.

Gastrointestinal: nausea, vomiting, epigastric distress, abdominal cramping, diarrhea, constipation, mesenteric arterial thrombosis, ischemic colitis.

Allergic: pharyngitis and agranulocytosis, erythematous rash, fever combined with aching and sore throat, laryngospasm and respiratory distress.

Respiratory: bronchospasm.

Hematologic: agranulocytosis, nonthrombocytopenic purpura, thrombocytopenic purpura.

Auto-Immune: In extremely rare instances, systemic lupus erythematosus has been reported.

Miscellaneous: alopecia, LE-like reactions, psoriasisiform rashes, dry eyes, male impotence, and Peyronie's disease have been reported rarely. Oculomucocutaneous reactions involving the skin, serous membranes and conjunctivae reported for a beta blocker (practolol) have not been associated with propranolol.

*The appearance of INDERAL tablets is a registered trademark of Ayerst Laboratories.

Copyright © 1985 Ayerst Laboratories

9429/185



AYERST LABORATORIES
New York, N.Y. 10017

Before prescribing, see complete prescribing information in SK&F CO. literature or PDR. The following is a brief summary.

*** WARNING**

This drug is not indicated for initial therapy of edema or hypertension. Edema or hypertension requires therapy titrated to the individual. If this combination represents the dosage so determined, its use may be more convenient in patient management. Treatment of hypertension and edema is not static, but must be reevaluated as conditions in each patient warrant.

Contraindications: Concomitant use with other potassium-sparing agents such as spironolactone or amiloride. Further use in anuria, progressive renal or hepatic dysfunction, hyperkalemia. Pre-existing elevated serum potassium. Hypersensitivity to either component or other sulfonamide-derived drugs.

Warnings: Do not use potassium supplements, dietary or otherwise, unless hypokalemia develops or dietary intake of potassium is markedly impaired. If supplementary potassium is needed, potassium tablets should not be used. Hyperkalemia can occur, and has been associated with cardiac irregularities. It is more likely in the severely ill, with urine volume less than one liter/day, the elderly and diabetics with suspected or confirmed renal insufficiency. Periodically, serum K⁺ levels should be determined. If hyperkalemia develops, substitute a thiazide alone, restrict K⁺ intake. Associated widened QRS complex or arrhythmia requires prompt additional therapy. Thiazides cross the placental barrier and appear in cord blood. Use in pregnancy requires weighing anticipated benefits against possible hazards, including fetal or neonatal jaundice, thrombocytopenia, other adverse reactions seen in adults. Thiazides appear and triamterene may appear in breast milk. If their use is essential, the patient should stop nursing. Adequate information on use in children is not available. Sensitivity reactions may occur in patients with or without a history of allergy or bronchial asthma. Possible exacerbation or activation of systemic lupus erythematosus has been reported with thiazide diuretics.

Precautions: The bioavailability of the hydrochlorothiazide component of 'Dyazide' is about 50% of the bioavailability of the single entity. Theoretically, a patient transferred from the single entities of Dyrenium (triamterene, SK&F CO.) and hydrochlorothiazide may show an increase in blood pressure or fluid retention. Similarly, it is also possible that the lesser hydrochlorothiazide bioavailability could lead to increased serum potassium levels. However, extensive clinical experience with 'Dyazide' suggests that these conditions have not been commonly observed in clinical practice. Do periodic serum electrolyte determinations (particularly important in patients vomiting excessively or receiving parenteral fluids, and during concurrent use with amphotericin B or corticosteroids or corticotropin [ACTH]). Periodic BUN and serum creatinine determinations should be made, especially in the elderly, diabetics or those with suspected or confirmed renal insufficiency. Cumulative effects of the drug may develop in patients with impaired renal function. Thiazides should be used with caution in patients with impaired hepatic function. They can precipitate coma in patients with severe liver disease. Observe regularly for possible blood dyscrasias, liver damage, other idiosyncratic reactions. Blood dyscrasias have been reported in patients receiving triamterene, and leukopenia, thrombocytopenia, agranulocytosis, and aplastic and hemolytic anemia have been reported with thiazides. Thiazides may cause manifestation of latent diabetes mellitus. The effects of oral anticoagulants may be decreased when used concurrently with hydrochlorothiazide; dosage adjustments may be necessary. Clinically insignificant reductions in arterial responsiveness to norepinephrine have been reported. Thiazides have also been shown to increase the paralyzing effect of nondepolarizing muscle relaxants such as tubocurarine. Triamterene is a weak folic acid antagonist. Do periodic blood studies in cirrhotics with splenomegaly. Antihypertensive effects may be enhanced in post-sympathectomy patients. Use cautiously in surgical patients. Triamterene has been found in renal stones in association with the other usual calculus components. Therefore, 'Dyazide' should be used with caution in patients with histories of stone formation. A few occurrences of acute renal failure have been reported in patients on 'Dyazide' when treated with indomethacin. Therefore, caution is advised in administering nonsteroidal anti-inflammatory agents with 'Dyazide'. The following may occur: transient elevated BUN or creatinine or both, hyperglycemia and glycosuria (diabetic insulin requirements may be altered), hyperuricemia and gout, digitalis intoxication (in hypokalemia), decreasing alkali reserve with possible metabolic acidosis. 'Dyazide' interferes with fluorescent measurement of quinidine. Hypokalemia is uncommon with 'Dyazide', but should it develop, corrective measures should be taken such as potassium supplementation or increased dietary intake of potassium-rich foods. Corrective measures should be instituted cautiously and serum potassium levels determined. Discontinue corrective measures and 'Dyazide' should laboratory values reveal elevated serum potassium. Chloride deficit may occur as well as dilutional hyponatremia. Concurrent use with chlorpropamide may increase the risk of severe hyponatremia. Serum PBI levels may decrease without signs of thyroid disturbance. Calcium excretion is decreased by thiazides. 'Dyazide' should be withdrawn before conducting tests for parathyroid function.

Thiazides may add to or potentiate the action of other antihypertensive drugs.

Diuretics reduce renal clearance of lithium and increase the risk of lithium toxicity.

Adverse Reactions: Muscle cramps, weakness, dizziness, headache, dry mouth; anaphylaxis, rash, urticaria, photosensitivity, purpura, other dermatological conditions; nausea and vomiting, diarrhea, constipation, other gastrointestinal disturbances; postural hypotension (may be aggravated by alcohol, barbiturates, or narcotics). Necrotizing vasculitis, paresthesias, icterus, pancreatitis, xanthopsia and respiratory distress including pneumonitis and pulmonary edema, transient blurred vision, sialadenitis, and vertigo have occurred with thiazides alone. Triamterene has been found in renal stones in association with other usual calculus components. Rare incidents of acute interstitial nephritis have been reported. Impotence has been reported in a few patients on 'Dyazide', although a causal relationship has not been established.

Supplied: 'Dyazide' is supplied as a red and white capsule, in bottles of 1000 capsules; Single Unit Packages (unit-dose) of 100 (intended for institutional use only); in Patient-Pak™ unit-of-use bottles of 100.

BRS-DZL39

a product of
SK&F CO.
Carolina, PR 00630

© SK&F Co., 1985

The unique
red and white
Dyazide® capsule:
Your assurance of
SK&F quality.



In Hypertensives Over 50, Diuretics Are Preferred

The 1984 Report of the Joint National Committee on Detection, Evaluation, and Treatment of High Blood Pressure recommends diuretics as the favored monotherapy in patients over 50 years of age, regardless of sex or race.



Beta Blockers Aren't for Everyone...
For Hypertensive Patients* Over 50

P R E S C R I B E[®]

25 mg Hydrochlorothiazide/50 mg Triamterene/SKF

Used with Confidence for over 19 Years

Serum K⁺ and BUN should be checked periodically (see Warnings and Precautions).

BALANCED CALCIUM CHANNEL BLOCKER



CARDIZEM[®]
(diltiazem HCl)

balances
potent
coronary
vasodilation
with a low
incidence of
side effects

Low incidence of side effects

CARDIZEM[®] (diltiazem HCl) produces an incidence of adverse reactions not greater than that reported with placebo therapy, thus contributing to the patient's sense of well-being.

Indicated in the treatment of angina pectoris due to coronary artery disease and in the management of chronic stable angina (exertion associated angina) in patients who cannot tolerate or who are refractory to beta-blockers and/or nitrates or who remain symptomatic on adequate doses of these agents.

Reduces angina attack frequency*

42% to 46% decrease reported in multicenter study.¹

Increases exercise tolerance*

In Bruce exercise test,² control patients averaged 8.0 minutes to onset of pain; Cardizem patients averaged 9.8 minutes ($P < .005$).

CARDIZEM[®]
(diltiazem HCl)

**THE BALANCED
CALCIUM CHANNEL BLOCKER**

Please see full prescribing information on following page.


2/84

PROFESSIONAL USE INFORMATION



DESCRIPTION

CARDIZEM® (diltiazem hydrochloride) is a calcium ion influx inhibitor (slow channel blocker or calcium antagonist). Chemically, diltiazem hydrochloride is 1,5-Benzothiazepin-4(5H)-one, 3-(acetyloxy)-5-[2-(dimethylamino)ethyl]-2,3-dihydro-2-(4-methoxyphenyl)-, monohydrochloride, (+)-cis-. The chemical structure is:



Diltiazem hydrochloride is a white to off-white crystalline powder with a bitter taste. It is soluble in water, methanol, and chloroform. It has a molecular weight of 450.98. Each tablet of CARDIZEM contains either 30 mg or 60 mg diltiazem hydrochloride for oral administration.

CLINICAL PHARMACOLOGY

The therapeutic benefits achieved with CARDIZEM are believed to be related to its ability to inhibit the influx of calcium ions during membrane depolarization of cardiac and vascular smooth muscle.

Mechanisms of Action. Although precise mechanisms of its antianginal actions are still being delineated, CARDIZEM is believed to act in the following ways:

1. Angina Due to Coronary Artery Spasm: CARDIZEM has been shown to be a potent dilator of coronary arteries both epicardial and subendocardial. Spontaneous and ergonovine-induced coronary artery spasm are inhibited by CARDIZEM.
2. Exertional Angina: CARDIZEM has been shown to produce increases in exercise tolerance, probably due to its ability to reduce myocardial oxygen demand. This is accomplished via reductions in heart rate and systemic blood pressure at submaximal and maximal exercise work loads.

In animal models, diltiazem interferes with the slow inward (depolarizing) current in excitable tissue. It causes excitation-contraction uncoupling in various myocardial tissues without changes in the configuration of the action potential. Diltiazem produces relaxation of coronary vascular smooth muscle and dilation of both large and small coronary arteries at drug levels which cause little or no negative inotropic effect. The resultant increases in coronary blood flow (epicardial and subendocardial) occur in ischemic and nonischemic models and are accompanied by dose-dependent decreases in systemic blood pressure and decreases in peripheral resistance.

Hemodynamic and Electrophysiologic Effects. Like other calcium antagonists, diltiazem decreases sinoatrial and atrioventricular conduction in isolated tissues and has a negative inotropic effect in isolated preparations. In the intact animal, prolongation of the AH interval can be seen at higher doses.

In man, diltiazem prevents spontaneous and ergonovine-provoked coronary artery spasm. It causes a decrease in peripheral vascular resistance and a modest fall in blood pressure and, in exercise tolerance studies in patients with ischemic heart disease, reduces the heart rate blood pressure product for any given work load. Studies to date, primarily in patients with good ventricular function, have not revealed evidence of a negative inotropic effect; cardiac output, ejection fraction, and left ventricular end diastolic pressure have not been affected. There are as yet few data on the interaction of diltiazem and beta-blockers. Resting heart rate is usually unchanged or slightly reduced by diltiazem.

Intravenous diltiazem in doses of 20 mg prolongs AH conduction time and AV node functional and effective refractory periods approximately 20%. In a study involving single oral doses of 300 mg of CARDIZEM in six normal volunteers, the average maximum PR prolongation was 14% with no instances of greater than first-degree AV block. Diltiazem-associated prolongation of the AH interval is not more pronounced in patients with first-degree heart block. In patients with sick sinus syndrome, diltiazem significantly prolongs sinus cycle length (up to 50% in some cases).

Chronic oral administration of CARDIZEM in doses of up to 240 mg/day has resulted in small increases in PR interval, but has not usually produced abnormal prolongation. There were, however, three instances of second-degree AV block and one instance of third-degree AV block in a group of 959 chronically treated patients.

Pharmacokinetics and Metabolism. Diltiazem is absorbed from the tablet formulation to about 80% of a reference capsule and is subject to an extensive first-pass effect, giving an absolute bioavailability (compared to intravenous dosing) of about 40%. CARDIZEM undergoes extensive hepatic metabolism in which 2% to 4% of the unchanged drug appears in the urine. In vitro binding studies show CARDIZEM is 70% to 80% bound to plasma proteins. Competitive ligand binding studies have also shown CARDIZEM binding is not altered by therapeutic concentrations of digoxin, hydrochlorothiazide, phenylbutazone, propranolol, salicylic acid, or warfarin. Single oral doses of 30 to 120 mg of CARDIZEM result in detectable plasma levels within 30 to 60 minutes and peak plasma levels two to three hours after drug administration. The plasma elimination half-life following single or multiple drug administration is approximately 3.5 hours. Desacetyl diltiazem is also present in the plasma at levels of 10% to 20% of the parent drug and is 25% to 50% as potent a coronary vasodilator as diltiazem. Therapeutic blood levels of CARDIZEM appear to be in the range of 50 to 200 ng/ml. There is a departure from dose-linearity when single doses above 60 mg are given; a 120-mg dose gave blood levels three times that of the 60-mg dose. There is no information about the effect of renal or hepatic impairment on excretion or metabolism of diltiazem.

INDICATIONS AND USAGE

1. Angina Pectoris Due to Coronary Artery Spasm. CARDIZEM

is indicated in the treatment of angina pectoris due to coronary artery spasm. CARDIZEM has been shown effective in the treatment of spontaneous coronary artery spasm presenting as Prinzmetal's variant angina (resting angina with ST-segment elevation occurring during attacks).

2. Chronic Stable Angina (Classic Effort-Associated Angina). CARDIZEM is indicated in the management of chronic stable angina. CARDIZEM has been effective in controlled trials in reducing angina frequency and increasing exercise tolerance.

There are no controlled studies of the effectiveness of the concomitant use of diltiazem and beta-blockers or of the safety of this combination in patients with impaired ventricular function or conduction abnormalities.

CONTRAINDICATIONS

CARDIZEM is contraindicated in (1) patients with sick sinus syndrome except in the presence of a functioning ventricular pacemaker, (2) patients with second- or third-degree AV block except in the presence of a functioning ventricular pacemaker, and (3) patients with hypotension (less than 90 mm Hg systolic).

WARNINGS

1. **Cardiac Conduction.** CARDIZEM prolongs AV node refractory periods without significantly prolonging sinus node recovery time, except in patients with sick sinus syndrome. This effect may rarely result in abnormally slow heart rates (particularly in patients with sick sinus syndrome) or second- or third-degree AV block (six of 1243 patients for 0.48%). Concomitant use of diltiazem with beta-blockers or digitalis may result in additive effects on cardiac conduction. A patient with Prinzmetal's angina developed periods of asystole (2 to 5 seconds) after a single dose of 60 mg of diltiazem.
2. **Congestive Heart Failure.** Although diltiazem has a negative inotropic effect in isolated animal tissue preparations, hemodynamic studies in humans with normal ventricular function have not shown a reduction in cardiac index nor consistent negative effects on contractility (dp/dt). Experience with the use of CARDIZEM alone or in combination with beta-blockers in patients with impaired ventricular function is very limited. Caution should be exercised when using the drug in such patients.
3. **Hypotension.** Decreases in blood pressure associated with CARDIZEM therapy may occasionally result in symptomatic hypotension.
4. **Acute Hepatic Injury.** In rare instances, patients receiving CARDIZEM have exhibited reversible acute hepatic injury as evidenced by moderate to extreme elevations of liver enzymes. (See PRECAUTIONS and ADVERSE REACTIONS.)

PRECAUTIONS

General. CARDIZEM (diltiazem hydrochloride) is extensively metabolized by the liver and excreted by the kidneys and in bile. As with any new drug given over prolonged periods, laboratory parameters should be monitored at regular intervals. The drug should be used with caution in patients with impaired renal or hepatic function. In subacute and chronic dog and rat studies designed to produce toxicity, high doses of diltiazem were associated with hepatic damage. In special subacute hepatic studies, oral doses of 125 mg/kg and higher in rats were associated with histological changes in the liver which were reversible when the drug was discontinued. In dogs, doses of 20 mg/kg were also associated with hepatic changes; however, these changes were reversible with continued dosing.

Drug Interaction. Pharmacologic studies indicate that there may be additive effects in prolonging AV conduction when using beta-blockers or digitalis concomitantly with CARDIZEM. (See WARNINGS.)

Controlled and uncontrolled domestic studies suggest that concomitant use of CARDIZEM and beta-blockers or digitalis is usually well tolerated. Available data are not sufficient, however, to predict the effects of concomitant treatment, particularly in patients with left ventricular dysfunction or cardiac conduction abnormalities. In healthy volunteers, diltiazem has been shown to increase serum digoxin levels up to 20%.

Carcinogenesis, Mutagenesis, Impairment of Fertility. A 24-month study in rats and a 21-month study in mice showed no evidence of carcinogenicity. There was also no mutagenic response in *in vitro* bacterial tests. No intrinsic effect on fertility was observed in rats.

Pregnancy. Category C. Reproduction studies have been conducted in mice, rats, and rabbits. Administration of doses ranging from five to ten times greater (on a mg/kg basis) than the daily recommended therapeutic dose has resulted in embryo and fetal lethality. These doses, in some studies, have been reported to cause skeletal abnormalities. In the perinatal/postnatal studies, there was some reduction in early individual pup weights and survival rates. There was an increased incidence of stillbirths at doses of 20 times the human dose or greater.

There are no well-controlled studies in pregnant women; therefore, use CARDIZEM in pregnant women only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers. It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, exercise caution when CARDIZEM is administered to a nursing woman if the drug's benefits are thought to outweigh its potential risks in this situation.

Pediatric Use. Safety and effectiveness in children have not been established.

ADVERSE REACTIONS

Serious adverse reactions have been rare in studies carried out to date, but it should be recognized that patients with impaired ventricular function and cardiac conduction abnormalities have usually been excluded.

In domestic placebo-controlled trials, the incidence of adverse reactions reported during CARDIZEM therapy was not greater than that reported during placebo therapy.

The following represent occurrences observed in clinical studies which can be at least reasonably associated with the pharmacology of calcium influx inhibition. In many cases, the relationship to CARDIZEM has not been established. The most common occurrences, as well as their frequency of presentation, are: edema (2.4%),

headache (2.1%), nausea (1.9%), dizziness (1.5%), rash (1.3%), asthenia (1.2%), AV block (1.1%). In addition, the following events were reported infrequently (less than 1%) with the order of presentation corresponding to the relative frequency of occurrence.

Cardiovascular:	Flushing, arrhythmia, hypotension, bradycardia, palpitations, congestive heart failure, syncope.
Nervous System:	Paresthesia, nervousness, somnolence, tremor, insomnia, hallucinations, and amnesia.
Gastrointestinal:	Constipation, dyspepsia, diarrhea, vomiting, mild elevations of alkaline phosphatase, SGOT, SGPT, and LDH.
Dermatologic:	Pruritus, petechiae, urticaria, photosensitivity.
Other:	Polyuria, nocturia.

The following additional experiences have been noted:

A patient with Prinzmetal's angina experiencing episodes of vasospastic angina developed periods of transient asymptomatic asystole approximately five hours after receiving a single 60-mg dose of CARDIZEM.

The following postmarketing events have been reported infrequently in patients receiving CARDIZEM: erythema multiforme; leukopenia; and extreme elevations of alkaline phosphatase, SGOT, SGPT, LDH, and CPK. However, a definitive cause and effect between these events and CARDIZEM therapy is yet to be established.

OVERDOSAGE OR EXAGGERATED RESPONSE

Overdosage experience with oral diltiazem has been limited. Single oral doses of 300 mg of CARDIZEM have been well tolerated by healthy volunteers. In the event of overdosage or exaggerated response, appropriate supportive measures should be employed in addition to gastric lavage. The following measures may be considered:

Bradycardia	Administer atropine (0.60 to 1.0 mg). If there is no response to vagal blockade, administer isoproterenol cautiously.
High-Degree AV Block	Treat as for bradycardia above. Fixed high-degree AV block should be treated with cardiac pacing.
Cardiac Failure	Administer inotropic agents (isoproterenol, dopamine, or dobutamine) and diuretics.
Hypotension	Vasopressors (eg, dopamine or levarterenol bitartrate).

Actual treatment and dosage should depend on the severity of the clinical situation and the judgment and experience of the treating physician.

The oral LD₅₀'s in mice and rats range from 415 to 740 mg/kg and from 560 to 810 mg/kg, respectively. The intravenous LD₅₀'s in these species were 60 and 38 mg/kg, respectively. The oral LD₅₀ in dogs is considered to be in excess of 50 mg/kg, while lethality was seen in monkeys at 360 mg/kg. The toxic dose in man is not known, but blood levels in excess of 800 ng/ml have not been associated with toxicity.

DOSE AND ADMINISTRATION

Exertional Angina Pectoris Due to Atherosclerotic Coronary Artery Disease or Angina Pectoris at Rest Due to Coronary Artery Spasm. Dosage must be adjusted to each patient's needs. Starting with 30 mg four times daily, before meals and at bedtime, dosage should be increased gradually (given in divided doses three or four times daily) at one- to two-day intervals until optimum response is obtained. Although individual patients may respond to any dosage level, the average optimum dosage range appears to be 180 to 240 mg/day. There are no available data concerning dosage requirements in patients with impaired renal or hepatic function. If the drug must be used in such patients, titration should be carried out with particular caution.

Concomitant Use With Other Antianginal Agents:

1. **Sublingual NITG** may be taken as required to abort acute anginal attacks during CARDIZEM therapy.
2. **Prophylactic Nitrate Therapy**—CARDIZEM may be safely administered with short- and long-acting nitrates, but there have been no controlled studies to evaluate the antianginal effectiveness of this combination.
3. **Beta-blockers.** (See WARNINGS and PRECAUTIONS.)

HOW SUPPLIED

Cardizem 30-mg tablets are supplied in bottles of 100 (NDC 0088-1771-47) and in Unit Dose Identification Paks of 100 (NDC 0088-1771-49). Each green tablet is engraved with MARION on one side and 1771 engraved on the other. CARDIZEM 60-mg scored tablets are supplied in bottles of 100 (NDC 0088-1772-47) and in Unit Dose Identification Paks of 100 (NDC 0088-1772-49). Each yellow tablet is engraved with MARION on one side and 1772 on the other.

Issued 4/1/84

Another patient benefit product from




In 7 of 10
anxious
geriatrics...

Anxiety and
depressive
symptoms
go hand
in hand.



Xanax rapidly relieves anxiety with depressive symptoms.



In a recent clinical study¹ of 83 geriatric patients with clinical anxiety, 73% were diagnosed as having symptoms of depressed mood.

XANAX is well suited for therapy because it demonstrates greater efficacy than placebo in reducing the Hamilton Anxiety Rating Scale Total Score and individual items including depressed mood (see Figure).

With clinical advantages for geriatric patients.

- **Rapidly relieves** the symptoms of anxiety
- **Rapidly relieves** associated depressed mood
- **Well tolerated**—mild, transient drowsiness, the most commonly reported side effect the first week of therapy, shows a marked decrease thereafter and is not significantly different from that of placebo
- **Does not cause** cardiotoxicity
- **Specific geriatric dosage**—0.25 mg, two or three times daily

1. Cohn JB: Double-blind safety and efficacy comparison of alprazolam and placebo in the treatment of anxiety in geriatric patients. *Curr Ther Res* 1984;35(1):100-112.



Xanax[®] 0.25 mg
Tablets
alprazolam[®]

©1984 The Upjohn Company

Upjohn THE UPJOHN COMPANY
Kalamazoo, Michigan 49001 USA

Please see next page for brief summary of prescribing information.

Classified Advertisements

The rate for each insertion is **\$5 per line** (average six words per line) with **five line (\$25) minimum**. Box number charge: \$5 each month.

Classified display rates \$50 per inch.

Copy for classified advertisements should be received not later than the **first of the month preceding issue**. All copy must be typed or printed. • Classified advertisers using Box Numbers forbid the disclosure of their identity. Your inquiries in writing will be forwarded to Box Number advertisers. The right is reserved to reject or modify all classified advertising copy in conformity with the decisions of the Advertising Committee.

Please Type or Print Advertising Copy

Classified Advertisements Are Payable in Advance

THE WESTERN JOURNAL OF MEDICINE
44 GOUGH STREET, SAN FRANCISCO, CA 94103

PHYSICIANS WANTED

WYOMING—ER STAFF POSITION—Emergency Room in Rawlins, WY with moderate volume. Independent contractor status. Malpractice paid. Very flexible schedule. Excellent hunting, fishing, back-packing and skiing in surrounding area. Send Résumé to: Duane Eugene Abels, DO PC, PO Box 1536, Rawlins, WY 82301 or call (307) 324-2221 x115.

FAMILY PRACTICE PHYSICIAN to join 15 physician multispecialty group located in Southern Idaho. Send résumé to Business Administrator, Box 1233, Twin Falls, ID 83301.

INSTITUTION PHYSICIAN needed to staff the Wyoming State Training School at Lander, Wyoming. We have 420 mentally retarded residents with a 30 bed Health Care Center with a staff of four physicians. Attractive salary and fringe benefits. No overhead. Pleasant surrounding area including an excellent medical group in Lander. If interested, please contact the Personnel Manager, Wyoming State Training School, 8204 State Highway 789, Lander, WY 82520. An Equal Opportunity Employer.

FAMILY PRACTICE opportunity in eastern New Mexico. Join two physicians in private practice situation with an established patient population. Must do OB. No buy-in; just share expenses and call. Family community with stable economy. Contact Bill Norris, Southwest Community Health Services, PO Box 26666, Albuquerque, NM 87125-6666, or call 1 (800) 545-4030, ext. 3050.

EUGENE, OREGON: FP/GP to join staff in salaried position with clinic providing outpatient and hospital care primarily to low income population. Half-time to start, flexible scheduling. Contact: Michael Weinstein, MD, 323 East 12th Ave, Eugene, OR 97401; (503) 484-4800.

ACADEMIC GENERAL INTERNISTS, CARDIOLOGIST, AND ENDOCRINOLOGIST, DO—Wanted for full-time faculty positions. Must be Board admissible or certified, interested in teaching medical students and house staff. Good balance between clinical practice and teaching with research opportunities available. Assistant Professor level with excellent guaranteed salary and benefits. Send CV to Victor H. Kaylarian, DO, Department of Medicine, University of Osteopathic Medicine and Health Sciences, 3200 Grand Ave., Des Moines, IA 50312. All inquiries confidential.

KETCHUM, SUN VALLEY, IDAHO: Internist, Surgeon, Pediatrician, Orthopedist in multispecialty, expanding, well-equipped clinic. Income moderate; quality of life highest. Dr Bryan Stone, PO Box 2198, Ketchum, ID 83340; (208) 726-9361.

PHYSICIANS WANTED

SOUTHERN CALIFORNIA

Prestigious HMO is seeking experienced specialists and general practitioners for our facilities in Los Angeles and Orange Counties. Located in close proximity to major teaching centers, we offer the opportunity for continued professional development and rewarding clinical practice. Excellent compensation and benefits including paid malpractice, life, disability, medical and dental coverage, paid vacations, sick leave, educational leave and retirement plan. Please send CV to: Director/Physician Recruitment, CIGNA Healthplans of California, 700 North Brand Blvd., Suite 500-49, Glendale, CA 91203.

OCCUPATIONAL MEDICINE STAFF POSITION available in well known Gulf South Multispecialty Clinic. Established full service occupational medicine program undergoing expansion. Excellent support from outstanding medical staff. Possible university appointment. Must be Board eligible/certified in Occupational Medicine with formal training in Occupational or Preventative Medicine. Excellent fringe benefits, salary negotiable. Send reply and CV to Box 6478, Western Journal of Medicine, 44 Gough St., San Francisco, CA 94103.

GENERAL SURGEON OR FAMILY PRACTITIONER with general surgery credentials wanted for a rural Washington State community. Full service hospital with drawing area of 6,000. Office space and ancillary services offered. Good opportunity in an agricultural four-season recreational area. Contact: Garvin Olson, Administrator, Janice Davis, Secretary, 1012 South Third, Dayton, WA 99328; (509) 382-2531.

Physicians

The Perfect Practice Is
One That Lets You Practice

At Westworld Community Healthcare, Inc., physicians have something very special: **Time**. Time to devote to their practices, and to their patients. And they have the time because we handle all the business details their practices demand.

The goal of every Westworld Hospital is to help our physicians be as productive as they can possibly be. We handle all the paperwork and administrative details, and assist in everything from setting up medical offices to providing staffing and equipment. What's more, we provide an excellent compensation package - competitive with private practice - along with incentive for additional income. And since our commitment is to meet the unique health care needs of rural America, there is no competing with hundreds of other practitioners for patients.

All of these advantages mean **security** for the Westworld physician. If your commitment is to the highest standards in patient care, and if a rural environment sounds inviting to you, investigate Westworld Community Healthcare. We have a variety of attractive practice opportunities in rural America. Call collect now for more information, or send your curriculum vitae to: **Mr. Joel Feinstein, Dept. PH-3, (714) 768-2981, Westworld Community Healthcare, Inc., 83832 Rockfield Rd., Lake Forest, CA 92630-2880.** We are an equal opportunity employer m/f.

WCH Westworld Community Healthcare, Inc.

HAYWARD, CALIFORNIA 20 man Bay Area multispecialty group with 4 pediatricians seeking Board certified or eligible Pediatrician to replace retired physician, May 1985. Group active in HMO's and satellite office. Contact: Donald L. Lass, Administrator, Hayward Medical Group, 27212 Calaroga Ave., Hayward, CA 94545; (415) 785-5000.

CALIFORNIA—Join a partnership of established physicians providing urgent and ambulatory care in both northern and southern California. Physicians earn a guaranteed minimum with a strong incentive package and have the opportunity to quickly become partners. Our attractive centers are rapidly growing! Please indicate your geographic preference (Los Angeles or San Francisco Bay Area) and contact: California Emergency Physicians, 440 Grand Ave., Suite 500, Oakland, CA 94610; (415) 832-6400.

(Continued on Page 136)

INDUSTRIAL MEDICAL CLINIC

SAN FRANCISCO BAY AREA — CLOSE TO SFO AIRPORT


The most dominant clinic in area. Central location amidst hundreds of employers in dense business/industry area. Extremely accessible. 100+ accounts and growing. First class facility in 7,200 sq. ft. suite with \$250,000 invested less than 3-years ago. Per its second complete year, Billings totalled \$708,000 with Collections being \$681,000. Focus to date has been work related injuries as Clinic referred out 300-orthopedic cases and 1,000 x-ray studies last year along with CT scans, pharmacy and inpatient/outpatient surgery. Management has not been able to capitalize on these built-in revenue areas, nor able to pursue thoughts on expanding into an Immediate Care Facility. Excellent opportunity for Buyer who has Management Team, working capital and marketing strategy to develop full potential. Full price \$500,000. Direct inquiry to San Francisco Office of PPS and request Prospectus.



Professional Practice Sales

Serving the Professions since 1966
Nationwide Services

SO. CALIFORNIA 364 E. First St., Tustin, CA 92680 (714) 832-0230
NO. CALIFORNIA 1428 Irving St., San Francisco, CA 94122 (415) 665-8500



The RU-TUSS[®] Family of Products

Liquid... Expectorant... with Hydrocodone... Tablets... Capsules



Boots Pharmaceuticals, Inc.
6540 LINE AVENUE, P.O. BOX 6750
SHREVEPORT, LOUISIANA 71106-9989

No matter what the season, you can depend on the RU-TUSS[®] Weather Service for a live, up-to-the-minute weather report anywhere in the United States. Contact your Boots Representative for toll-free access.

PHYSICIANS WANTED

PHYSICIAN WANTED: San Francisco B/C or B/E Internist/FP to join growing three man primary care group in San Francisco's Mission District. For further information contact Dr John Pierce, (415) 826-7575 or send résumé to Golden Gate Medical Group, 3620 Army St., San Francisco, CA 94110.

INFECTIOUS DISEASE AND INTERNAL MEDICINE—The Division of General Internal Medicine/Primary Care at Cook County Hospital wishes to recruit an Attending Physician to supervise care of both patients with general medical problems and patients with AIDS. Board eligibility in Internal Medicine is required; subspecialty training in Infectious Disease or Oncology is desirable. General Medicine responsibilities occupy 50 percent time and include ward attending and supervising the General Medicine Clinic. AIDS service responsibilities include participation in biweekly clinics and regular ward consultations for patients with AIDS and AIDS related conditions, and involvement in educational and research programs of the AIDS service. Sensitivity to the health care needs of gay men and other AIDS risk group members is needed. Starting salary for Board certified physicians is \$54,228 a year. For further information please contact Renslow Sherer, MD, Coordinator of AIDS Services, Division of General Medicine, Cook County Hospital, 1835 W. Harrison, Chicago, IL 60612. Cook County Hospital is an Equal Opportunity/Affirmative Action Employer.

INTERNIST-RHEUMATOLOGIST-SUBSPECIALIST. Opportunity to join solo Internist-Rheumatologist in private practice. Association and investment opportunity. Laboratory and x-ray. Send CV to R. S. Gordon, MD, 5945 Brockton Ave., Riverside, CA 92506.

WASHINGTON—Ambulatory Care Center, Tacoma. Full- or possibly part-time for FP/IM/EM trained physician. Contact Roger Simms, MD, (206) 759-6655 or reply with CV to 5702 North 26th, Tacoma, WA 98407.

HEMATOLOGIST-ONCOLOGIST to join Oncologist in progressive multispecialty group practice located in suburban San Francisco area. University affiliation and NCOG protocol participation encouraged. Competitive salary and benefits leading to partnership. Send résumé to Box 6479, The Western Journal of Medicine, 44 Gough St., San Francisco, CA 94103.

FAMILY PRACTICE-INTERNAL MEDICINE-SURGEON. Multispecialty clinic in Northern Idaho Panhandle. Nominal lease includes fully-equipped office and treatment suites with receptionist and janitorial services provided. Clinic includes lab, x-ray and pharmacy. Billing services available. Located in a four-season playground rich with forests, lakes, streams and wild game. Just 20 minutes from a 14-run ski resort. Attractive lifestyle. For information: (208) 784-1221, ext. 304. Mail résumé to: Shoshone Medical Center, Jacobs Gulch, Kellogg, ID 83837.

IMMEDIATE OPENING—Opportunity for Internist, Cardiologist or Family Practice to join two doctor clinic grossing near \$400,000. Opportunity to buy in also. Two Surgeons and over five Family Practice Doctors who will support an Internist. Central WA; (509) 422-2600.

MEDICAL DIRECTOR—Division of Industrial Accidents, provides medical expertise for the staff of the Workers' Compensation Appeals Board; reviews work injury cases; coordinates various committees; acts as liaison for the WCAB and the medical and workers' compensation communities. Experience with industrial injuries required for this diverse and challenging civil service position. Good salary with excellent state benefits. Contact William L. Clark, MD, D/A Medical Bureau or Mike Giles, Director Personnel, 525 Golden Gate Ave., San Francisco, CA 94102; (415) 557-2370 or 557-3362. Final filing date 8-2-85. Equal opportunity employer.

PHYSICIANS WANTED

FAMILY PRACTITIONER—OUTPATIENT CARE: Full time position available for residency trained, Board Eligible/Board Certified Family Practitioner interested in a position involving outpatient care. Pediatrics, Prenatal Care, Adult Medicine, Orthopedics, and minor surgery responsibilities are available. For information, call Donald B. Dawson, Director of Physician Staffing, toll-free at 1 (800) 446-2255 [in California call 1 (800) 336-2255]. FHP Professional Staffing, 400 Oceangate Blvd., Suite 1317, Long Beach, CA 90802. For opportunities in Utah, call Maryalys Poulson collect at (801) 355-1234.

FAMILY PRACTITIONER—Full time position available for residency trained, Board Eligible/Board Certified Family Practitioners interested in practicing in a comprehensive care environment. Outpatient care and in-hospital responsibilities are offered in a growing family practice organization. Administrative opportunities also available. For information, call Donald B. Dawson, Director of Physician Staffing, toll-free at 1 (800) 446-2255 [in California call 1 (800) 336-2255]. FHP Professional Staffing, 400 Oceangate Blvd., Suite 1317, Long Beach, CA 90802. For opportunities in Utah, call Maryalys Poulson collect at (801) 355-1234.

CALIFORNIA, SAN JOSE: Emergency Physician sought for position as part-time Director of Poison Control Center at a university-affiliated teaching hospital. Responsibilities include patient care in the Emergency Department. Must be Board Prepared/Board Certified in Clinical Toxicology. Fee-for-service with minimum guarantee. Contact: James B. Lane, MD, 1625 The Alameda, #201, San Jose, CA 95126; (408) 293-8881.

CALIFORNIA—Board Certified/eligible or residency trained emergency physician wanted to join 320 physician multispecialty group. Competitive salary and excellent fringe benefits. Ninety miles from Sierra skiing and San Francisco. California license required. Send curriculum vitae to Mrs. Carolyn Whelan, The Permanente Medical Group, Inc., PO Box 254999, Sacramento, CA 95825. An equal opportunity employer.

GENERAL INTERNIST—BC/BE to join four man group of internists located in beautiful central coastal community. Senior partner retiring January 1986. Excellent opportunity for young energetic physician; salary plus percentage if mutually satisfactory. Write: Box 6482, Western Journal of Medicine, 44 Gough St., San Francisco, CA 94103.

PATHOLOGIST: Large pathology group seeking qualified anatomic and clinical pathologist. Excellent position with growth potential. Send résumé to: Glenview Pathology Medical Group, 9331 Venice Blvd., Culver City, CA 90230.

PHYSICIANS WANTED

TIBURON, CALIFORNIA: Senior Internist seeking Internist or other specialist willing to share small, well-equipped and well-staffed office, one to three days each week. For terms and particulars call Mrs. Grossi, (415) 435-0656. Business hours 9:30-5:00.

FAMILY PRACTITIONER—CALIFORNIA CLINIC: Small group, hospital practice and shared call system. Send CV or call John Aleman, MD, Box 858, Merced, CA 95341; (209) 383-1848.

PRIMARY CARE/URGENT CARE: Primary Care—Northern and Southern California: Part- or full-time positions for Primary Care physicians experienced in treatment of industrial type trauma for Occupational Health Clinics. Fringe benefits, malpractice insurance provided.

Urgent Care—Northern and Southern California: Rapidly expanding medical group servicing Urgent Care network throughout California (San Diego, Orange County, San Joaquin Valley and Bay areas) needs independent contractors with primary care, ER, occupational medicine specialties. California license required, prior out-patient practice a plus. Client provides all personnel, facility, ancillary equipment with group providing malpractice. Guaranteed minimum plus monthly incentive, stock options available. No relocation assistance. Write: Personnel, Urgent Care Centers of America, Inc., 2737 Walsh Ave., Santa Clara, CA 95051; (408) 727-1900. EOE.

THROUGHOUT CALIFORNIA: Immediate full- or part-time opening with California's leading provider of industrial/private care. Orthopedist and Neurologist needed for consultations, out-patient minor trauma. Industrial/Urgent Care Clinic. No nights, weekends, salary guarantee plus high incentive bonus. Malpractice and fringe benefits. Write Personnel, Urgent Care Centers of America, Inc., 2737 Walsh Ave., Santa Clara, CA 95051.

INTERNAL MEDICINE: Excellent opportunity with California's established leading provider of executive fitness/industrial care, for Internist (Board Certified/Eligible). Immediate full time/part time opening. Salary guaranteed plus production bonus. Malpractice and fringe benefits. Write Personnel, Urgent Care Centers of America, Inc., 2737 Walsh Ave., Santa Clara, CA 95051; (408) 727-1900. EOE.

FAMILY PRACTICE at scenic Lake Chelan, WA. Join 3 other BC FPs, share call, expenses and building. Obstetrics required. Nearly new hospital. Gross 180-200K. Extensive recreational opportunities available. Practice for sale for modest negotiable amount or possibly available on a salaried basis. Contact Lake Chelan Clinic, PO Box 368, Chelan, WA 98816; (509) 682-2511.

(Continued on Page 138)



SPECTRUM EMERGENCY CARE, Inc.

Career emergency medicine positions are available with the nation's largest group in the following states:

California
Colorado
Idaho
Kansas

Montana
Nebraska
Nevada
N. Dakota

S. Dakota
Utah
Washington
Wyoming

Spectrum provides career emergency physicians with a competitive income; professional liability insurance; and reimbursement of CME tuition, ACEP dues, ACLS and ATLS training. For complete details on positions available in the states listed above contact:

Jan Bird
Spectrum Emergency Care, Inc.
6275 Lehman Drive
Suite C 202
Colorado Springs, CO 80918

1-800-525-3681 / 303-590-1755 / 1-800-421-6655

ARE YOU A NATURAL FOR AMWA MEMBERSHIP?

YOU ARE if you are a physician, a nurse, or a pharmacist who wants to submit a readable and publishable article to a professional journal.

YOU ARE if you are a professional writer, an editor, or a graphics designer in the health care field who wants to become even more creatively productive.

YOU ARE if you are a medical communicator who wants to know and use the best contemporary verbal/visual techniques for conveying information.

YOU ARE if you want to meet, on social and professional levels, the many kinds of people described who conduct and participate in relevant, stimulating workshops and seminars in all parts of the country.

YOU ARE if you fill out, clip, and mail this coupon.



Executive Secretary, AMWA
5272 River Road, Suite 290
Bethesda, Maryland 20016

YES. I want to know more about AMWA.

NAME _____

TITLE (or specialty) _____

ADDRESS _____

CITY _____ STATE _____ ZIP _____

AMERICAN
MEDICAL WRITERS
ASSOCIATION

NORTHERN CALIFORNIA: Physician, Board certified/eligible in Emergency Medicine or Family Practice, to work part-time or full-time in busy free-standing emergency center. Competitive compensation and malpractice paid. Send CV and references to Joanne Berkowitz, MD, The Emergency Center Medical Group, 4948 San Juan Ave., Fair Oaks, CA 95628.

FAMILY PRACTITIONER—TULARE COUNTY, CALIFORNIA: Board certified/eligible Family Practitioner to practice in an out-patient clinic, which includes in-patient duties, with 13 physicians. Consider a semi-rural lifestyle with cultural amenities, the Coast and the Sierra Nevada Mountains easily accessed. Salary: \$81,236-\$85,376 annually. The County provides a benefit package which includes malpractice insurance coverage. Send CV to Tulare County Personnel, Courthouse, Room 106, Visalia, CA 93291; (209) 733-6266. An Affirmative Action Employer.

CALIFORNIA—Physicians interested in fulltime practice in Urgent Care Center at Palo Alto Medical Foundation. Must be residency trained and Board Certified in FP, IM, General Surgery, Occupational Medicine or Emergency Medicine. ACLS Certified. Must be experienced in a wide variety of out-patient Medical/Surgical skills in adults and children. Occupational Medicine experience desirable. Salary, vacation, meeting time; malpractice, dental, medical, disability insurance; three year track for partnership. Available July/August 1985. For information send CV to: William E. Straw, MD, Medical Director, Urgent Care Center, Palo Alto Medical Foundation, 300 Homer Ave., Palo Alto, CA 94301. Please reply Box 6484, Western Journal of Medicine, 44 Gough St., San Francisco, CA 94103.

CALIFORNIA, SACRAMENTO: Partnership opportunity with established multi-hospital group practicing in Northern California. Full-time positions available for board-prepared or board-certified emergency physicians. Competitive salary and benefits. Malpractice paid. All hospitals with moderate volumes, many act as EMS base-stations. Send CV to Sacramento Emergency Medical Group, PO Box 214584, Sacramento, CA 95821.

WASHINGTON COASTAL COMMUNITY serving a population of 65,000 is actively recruiting the following specialists: ENT, Urologist, Orthopedic Surgeon, General Surgeon. A variety of practice support options are available, i.e., office space, relocation assistance, etc. Enjoy the support of major West Coast Catholic Hospital System. Community has close proximity to major recreational areas and easy access to Seattle and Portland. For information send CV and references to: Nancy Friedrich, The Friedrich Group, 9284 Ferncliff North East, Bainbridge Island, WA 98110.

CALIFORNIA, RURAL AMERICAN AND OVERSEAS: Primary care physicians and OB/GYN needed for locum and permanent placements in CA, Saudi Arabia and, southwestern US. Excellent financial package, practice management and affiliation with a dynamic healthcare company. CV to: Beverly Frole, Westworld Healthcare Resources, 23832 Rockfield Rd., Lake Forest, CA 92630; (800) 847-1596.

MINOR EMERGENCY CENTERS located along the beautiful Wasatch front in Utah looking for qualified physicians. Send CV to Val Rollins, MD, Emergency Department, St. Benedict's Hospital, 5475 South 500, East Ogden, UT 84405.

URGENT CARE: Full- and part-time positions for physicians to staff Urgent Care Department of multispecialty group located 30 miles south of Santa Barbara, CA. Compensation—salary, hourly contract rate or percentage. Send résumé to Physician Recruitment, 2705 Loma Vista Rd., Ventura, CA 93003.

WESTERN U.S. OPENINGS

Several multi-specialty groups and clinics have asked us to recruit for over 100 positions of various specialties.

Western States Physician Search
240 West Shaw, Suite C
Clovis, CA 93612
(209) 297-7748

PHYSICIANS WANTED

CHAIRPERSON, DEPARTMENT OF SURGERY: The Permanente Medical Group is taking applications for the position of chairperson for the Department of Surgery at our Oakland Medical Center. The Oakland facility is a 330-bed acute care teaching hospital with a five-year general surgery residency program. The Surgery Department is composed of seven board certified general surgeons; two having completed vascular fellowships. Interested candidates must be board certified and prior experience with resident training programs preferred. Please send curriculum vitae to: Richard Brown, MD, Chairman, Search Committee, Kaiser Permanente Medical Group, Inc., 280 West MacArthur Blvd., Oakland, CA 94611.

FAMILY PRACTICE—MARIN COUNTY, CALIFORNIA: Rare opportunity to share office space and call in a community based private practice located one hour north of San Francisco. Interest in women's health care a plus. Box 255, Fairfax, CA 94930; (415) 488-4525 (evenings).

CARDIOLOGIST: To join two busy invasive Cardiologists in the beautiful Pacific Northwest. Practice includes open heart surgery program, PTCA. Requires catheterization and all non-invasive modalities. EPS and PTCA would be a plus. Please reply with a CV to Box 6487, Western Journal of Medicine, 44 Gough St., San Francisco, CA 94103.

URGENT CARE CLINIC: Sacramento, CA. Full time family practice oriented emergency physician. Twelve hours/day. Guarantee plus percentage. Malpractice provided. Minimum one year commitment. Please reply to Box 6485, Western Journal of Medicine, 44 Gough St., San Francisco, CA 94103.

MONTANA FAMILY PRACTICE: Have the best of both worlds. BE/BC Family Practitioner needed to establish a Department of Family Practice as part of The Billings Clinic—a 62-physician multispecialty group. Enjoy practice with or without obstetrics in a regional medical center with subspecialty backup and the cultural advantages of good schools, art, drama, symphony orchestra, etc. You and your family can enjoy the wide open spaces together. There is riding, fishing, hiking, skiing, hunting, canoeing, to name a few. This is a delightful place to raise a family. Salary leading to partnership. Contact: Paul V. Hoyer, MD, The Billings Clinic, PO Box 35100, Billings, MT 59107-5100. Telephone (406) 256-2500.

FAMILY PHYSICIANS WANTED: Openings for two Family Physicians in south central Idaho. Rural practice with specialists/Regional Medical Center 40 miles away. Excellent outdoor recreation. \$45,000-\$48,000, paid malpractice, CME leave, stipend, three day weekend twice a month. Contact Cookie Atkins, Family Health Services, PO Box 731, Twin Falls, ID 83303; (208) 734-3312.

GENERAL/ORTHOPEDIC SURGEON/FP-INTERNIST to join multispecialty group in New Mexico. Excellent benefit package. First year salary guarantee negotiable. Well-equipped private outpatient clinic. Five minutes from new hospital. Call Mr. Miller or write 114 West 11th St., Silver City, NM 88061; (505) 388-1511, or (505) 538-2408 in evenings.

MEDICAL OFFICE CONDOMINIUM SUITE FOR SALE OR LEASE: Hilltop Professional Building, 1855 San Miguel Dr., Walnut Creek, CA; park-like setting; ample parking; for information call App and Edwards Company at (415) 935-6970.



THOMAS WEBBER
General Director

TALK TO PATIENTS IN EVERYDAY SPANISH

Practical cassettes, native speakers, English explanations; software; courses/homestays in Mexico. Complete tape courses available from basic to advanced.

- Mexican Spanish Pronunciation/Making Friends in Spanish (Cassette & illustrated manual): \$13.95.
- Practical Medical Spanish (6-Cassette Course): \$83.45. California residents add 6% tax.
- SATISFACTION GUARANTEED.

SUMMER SCHOOL IN BAJA CALIFORNIA: Write for brochures on courses in Mexico, tapes, and software. California Spanish Language Association, 189-J Correo Fronterizo, San Ysidro, CA 92073.

LOCUM TENENS

LOCUM TENENS SERVICE WESTERN PHYSICIANS REGISTRY

... offers coverage for vacation or continuing education. To arrange coverage for your practice or to participate as temporary physician, contact: Carol Sweig, Director, 1124 Bal-lena, Alameda, CA 94501; (415) 521-4110.

PRACTICE AVAILABLE

OPHTHALMOLOGY PRACTICE FOR SALE: 1985 or 1986. Pacific Northwest Coast. Office building for lease or sale. Retiring. Will stay to introduce. Reply including CV to Box 6472, Western Journal of Medicine, 44 Gough St., San Francisco, CA 94103.

CALIFORNIA: Otolaryngology, Pediatric, Psychiatric, Ophthalmology, Allergy, OBG, Family, Internal, Surgery, Orthopedic, others. Contact Mary Bradshaw, Practice Broker/Recruiter, 21 Altamont Dr., Orinda, CA 94563; (415) 376-0762.

PULMONARY DISEASE/ALLERGY practice in San Bernardino. Good lease, 1,400-foot office. Computer billing and Video patient educational systems, Pulmonary Function Laboratory equipment, trained assistant. Donald Thornton, MD, 1800 North Western Ave., 3406, San Bernardino, CA 92411; (714) 887-8088.

WELL ESTABLISHED GENERAL PRACTICE and free standing care center in Northern California for sale. Office fully equipped. Inquiries Box 6481, Western Journal of Medicine, 44 Gough St, San Francisco, CA 94103.

WOMAN FP RETIRING: Will sell practice and equipment to FP/IM. Prime location. Terms negotiable. Box 26852, Albuquerque, NM 87125.

INTERNIST/FAMILY PRACTITIONER to take over existing practice in Los Angeles. Japanese bilingual physician would be perfect candidate: Board Eligible or Certified preferred. Truly once in a life-time opportunity. Please forward CV to Box 6483, Western Journal of Medicine, 44 Gough St., San Francisco, CA 94103.

CONTRACT NEGOTIATION

Practicing physicians with ten years' experience in collective bargaining and corporate medical affairs. Non-union. Call (415) 461-2080 for further information or write to PMC, Box 1434, Ross, CA 94957. Reply in strictest confidence.

PATIENT NEWSLETTER

Keeping in touch shows that you care! Send for your free sample copy of our quarterly health promotion newsletter. Customized for you. Mailing service option. Rx: Live well, 6809 Santa Maria Ln., Dallas, TX 75214-2847 or call (214) 821-5218.

MID-LEVEL PRACTITIONERS NATIONWIDE are seeking positions with Physicians, Groups, Clinics, and hospitals. We match your needs with theirs, before you meet! Save time, energy, money. MEDICAL INTERFACE, 18901 61st Ave. NE, Seattle, WA 98155; (206) 486-2814.

ALL MEDICAL AND SURGICAL SPECIALTIES sought by quality oriented, nationwide, referral service for medical-legal consultation. Highly rewarding opportunity for qualified experts to provide objective reviews. Specific terms strictly between participating expert and requesting party. Send inquiries to ENET 2020 West State St., Suite 164, Milwaukee, WI 53223.

PROFESSIONAL RÉSUMÉ SERVICES: Curriculum vitae preparation for physicians. Prompt and confidential. Call toll free. 1 (800) 6-CAREER. In PA, (215) 433-4112. Or write to PO Box AG, Allentown, PA 18106.

RADIOLOGIST SEEKS POSITION—41-year-old experienced Radiologist (academics and private practice) seeks multispecialty group, solo-type practice, or imaging center. Reply Box 6486, Western Journal of Medicine, 44 Gough St., San Francisco, CA 94103.

“When the Ayerst rep told me
it costs about 45¢ a day,
I said you can stop right there.”



Most doctors are pleasantly surprised to learn that the average cost of daily therapy with the world's most widely used beta blocker is so little, not much more than the cost of a daily newspaper.

When it's **INDERAL** (propranolol hydrochloride) tablets you want for your hypertension patients, remember to specify Dispense As Written (DAW) or Do Not Substitute on your prescriptions. That way, you can always be assured they'll get **INDERAL**®. Please see next page for brief summary of prescribing information.

INDERAL®
(PROPRANOLOL HCl)
Small price to pay.

“When the Ayerst rep told me
it costs about 45¢ a day,
I said you can stop right there.”

INDERAL[®] TABLETS
(PROPRANOLOL HCl)



BRIEF SUMMARY (FOR FULL PRESCRIBING INFORMATION, SEE PACKAGE CIRCULAR.)

INDERAL[®] (propranolol hydrochloride) Tablets

CONTRAINDICATIONS

INDERAL is contraindicated in 1) cardiogenic shock, 2) sinus bradycardia and greater than first degree block, 3) bronchial asthma, 4) congestive heart failure (see WARNINGS) unless the failure is secondary to a tachyarrhythmia treatable with INDERAL.

WARNINGS

CARDIAC FAILURE: Sympathetic stimulation may be a vital component supporting circulatory function in patients with congestive heart failure, and its inhibition by beta blockade may precipitate more severe failure. Although beta blockers should be avoided in overt congestive heart failure, if necessary they can be used with close follow-up in patients with a history of failure who are well compensated and are receiving digitalis and diuretics. Beta-adrenergic blocking agents do not abolish the inotropic action of digitalis on heart muscle.

IN PATIENTS WITHOUT A HISTORY OF HEART FAILURE: continued use of beta blockers can, in some cases, lead to cardiac failure. Therefore, at the first sign or symptom of heart failure, the patient should be digitalized and/or treated with diuretics, and the response observed closely, or INDERAL should be discontinued (gradually, if possible).

IN PATIENTS WITH ANGINA PECTORIS: there have been reports of exacerbation of angina and, in some cases, myocardial infarction, following abrupt discontinuance of INDERAL therapy. Therefore, when discontinuance of INDERAL is planned the dosage should be gradually reduced over at least a few weeks and the patient should be cautioned against interruption or cessation of therapy without the physician's advice. If INDERAL therapy is interrupted and exacerbation of angina occurs, it usually is advisable to reinstitute INDERAL therapy and take other measures appropriate for the management of unstable angina pectoris. Since coronary artery disease may be unrecognized, it may be prudent to follow the above advice in patients considered at risk of having occult atherosclerotic heart disease who are given propranolol for other indications.

Nonallergic Bronchospasm (e.g., chronic bronchitis, emphysema)— PATIENTS WITH BRONCHOSPASTIC DISEASES SHOULD IN GENERAL NOT RECEIVE BETA BLOCKERS. INDERAL should be administered with caution since it may block bronchodilation produced by endogenous and exogenous catecholamine stimulation of beta receptors.

MAJOR SURGERY: The necessity or desirability of withdrawal of beta-blocking therapy prior to major surgery is controversial. It should be noted, however, that the impaired ability of the heart to respond to reflex adrenergic stimuli may augment the risks of general anesthesia and surgical procedures.

INDERAL, like other beta blockers, is a competitive inhibitor of beta-receptor agonists and its effects can be reversed by administration of such agents, e.g., dobutamine or isoproterenol. However, such patients may be subject to protracted severe hypotension. Difficulty in starting and maintaining the heartbeat has also been reported with beta blockers.

DIABETES AND HYPOLYCEMIA: Beta-adrenergic blockade may prevent the appearance of certain premonitory signs and symptoms (pulse rate and pressure changes) of acute hypoglycemia in labile insulin-dependent diabetes. In these patients, it may be more difficult to adjust the dosage of insulin.

THYROTOXICOSIS: Beta blockade may mask certain clinical signs of hyperthyroidism. Therefore, abrupt withdrawal of propranolol may be followed by an exacerbation of symptoms of hyperthyroidism, including thyroid storm. Propranolol does not distort thyroid function tests.

IN PATIENTS WITH WOLFF-PARKINSON-WHITE SYNDROME, several cases have been reported in which, after propranolol, the tachycardia was replaced by a severe bradycardia requiring a demand pacemaker. In one case this resulted after an initial dose of 5 mg propranolol.

PRECAUTIONS

General: Propranolol should be used with caution in patients with impaired hepatic or renal function. INDERAL is not indicated for the treatment of hypertensive emergencies.

Beta-adrenoreceptor blockade can cause reduction of intraocular pressure. Patients should be told that INDERAL (propranolol hydrochloride) may interfere with the glaucoma screening test. Withdrawal may lead to a return of increased intraocular pressure.

Clinical Laboratory Tests: Elevated blood urea levels in patients with severe heart disease, elevated serum transaminase, alkaline phosphatase, lactate dehydrogenase.

DRUG INTERACTIONS: Patients receiving catecholamine-depleting drugs such as reserpine should be closely observed if INDERAL is administered. The added catecholamine-blocking action may produce an excessive reduction of resting sympathetic nervous activity which may result in hypotension, marked bradycardia, vertigo, syncopal attacks, or orthostatic hypotension.

Carcinogenesis, Mutagenesis, Impairment of Fertility: Long-term studies in animals have been conducted to evaluate toxic effects and carcinogenic potential. In 18-month studies in both rats and mice, employing doses up to 150 mg/kg/day, there was no evidence of significant drug-induced toxicity. There were no drug-related tumorigenic effects at any of the dosage levels. Reproductive studies in animals did not show any impairment of fertility that was attributable to the drug.

Pregnancy: Pregnancy Category C. INDERAL has been shown to be embryotoxic in animal studies at doses about 10 times greater than the maximum recommended human dose.

There are no adequate and well-controlled studies in pregnant women. INDERAL should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers: INDERAL is excreted in human milk. Caution should be exercised when INDERAL is administered to a nursing woman.

Pediatric Use: Safety and effectiveness in children have not been established.

ADVERSE REACTIONS

Most adverse effects have been mild and transient and have rarely required the withdrawal of therapy.

Cardiovascular: bradycardia; congestive heart failure; intensification of AV block; hypotension; paresthesia of hands; thrombocytopenic purpura; arterial insufficiency, usually of the Raynaud type.

Central Nervous System: Lightheadedness; mental depression manifested by insomnia, lassitude, weakness, fatigue; reversible mental depression progressing to catatonia; visual disturbances; hallucinations; an acute reversible syndrome characterized by disorientation for time and place, short-term memory loss, emotional lability, slightly clouded sensorium, and decreased performance on neuropsychometrics.

Gastrointestinal: nausea, vomiting, epigastric distress, abdominal cramping, diarrhea, constipation, mesenteric arterial thrombosis, ischemic colitis.

Allergic: pharyngitis and agranulocytosis, erythematous rash, fever combined with aching and sore throat, laryngospasm and respiratory distress.

Respiratory: bronchospasm.

Hematologic: agranulocytosis, nonthrombocytopenic purpura, thrombocytopenic purpura.

Auto-immune: In extremely rare instances, systemic lupus erythematosus has been reported.

Miscellaneous: alopecia, LE-like reactions, psoriasisiform rashes, dry eyes, male impotence, and Peyronie's disease have been reported rarely. Oculomucocutaneous reactions involving the skin, serous membranes and conjunctivae reported for a beta blocker (practolol) have not been associated with propranolol.

*The appearance of INDERAL tablets is a registered trademark of Ayerst Laboratories.

Copyright © 1985 Ayerst Laboratories

9418/185

Ayerst

AYERST LABORATORIES
New York, N.Y. 10017

New Prescribing Information

Now—Easier to remember...
easier to prescribe

In place of Limbitrol 5-12.5 write:

Limbitrol[®]
Each tablet contains 6 mg clordiazepoxide and 12.5 mg amitriptyline (as the hydrochloride salt).

In place of Limbitrol 10-25 write:

Limbitrol DS[®]
Each tablet contains 10 mg clordiazepoxide and 25 mg amitriptyline (as the hydrochloride salt).

Before prescribing, please consult complete product information, a summary of which follows:

Indications: Relief of moderate to severe depression associated with moderate to severe anxiety.

Contraindications: Known hypersensitivity to benzodiazepines or tricyclic antidepressants. Do not use with monoamine oxidase (MAO) inhibitors or within 14 days following discontinuation of MAO inhibitors since hyperpyretic crises, severe convulsions and deaths have occurred with concomitant use, then initiate cautiously, gradually increasing dosage until optimal response is achieved. Contraindicated during acute recovery phase following myocardial infarction.

Warnings: Use with great care in patients with history of urinary retention or angle-closure glaucoma. Severe constipation may occur in patients taking tricyclic antidepressants and anticholinergic-type drugs. Closely supervise cardiovascular patients. (Arrhythmias, sinus tachycardia and prolongation of conduction time reported with use of tricyclic antidepressants, especially high doses. Myocardial infarction and stroke reported with use of this class of drugs.) Caution patients about possible combined effects with alcohol and other CNS depressants and against hazardous occupations requiring complete mental alertness (e.g., operating machinery, driving).

Usage in Pregnancy: Use of minor tranquilizers during the first trimester should almost always be avoided because of increased risk of congenital malformations as suggested in several studies. Consider possibility of pregnancy when instituting therapy; advise patients to discuss therapy if they intend to or do become pregnant.

Since physical and psychological dependence to clordiazepoxide have been reported rarely, use caution in administering Limbitrol to addiction-prone individuals or those who might increase dosage, withdrawal symptoms following discontinuation of either component alone have been reported (nausea, headache and malaise for amitriptyline; symptoms [including convulsions] similar to those of barbiturate withdrawal for clordiazepoxide).

Precautions: Use with caution in patients with a history of seizures, in hyperthyroid patients or those on thyroid medica-

tion, and in patients with impaired renal or hepatic function. Because of the possibility of suicide in depressed patients, do not permit easy access to large quantities in these patients.

Periodic liver function tests and blood counts are recommended during prolonged treatment. Amitriptyline component may block action of guanethidine or similar antihypertensives. Concomitant use with other psychotropic drugs has not been evaluated; sedative effects may be additive. Discontinue several days before surgery. Limit concomitant administration of ECT to essential treatment. See Warnings for precautions about pregnancy. Limbitrol should not be taken during the nursing period. Not recommended in children under 12. In the elderly and debilitated, limit to smallest effective dosage to preclude ataxia, oversedation, confusion or anticholinergic effects.

Adverse Reactions: Most frequently reported are those associated with either component alone: drowsiness, dry mouth, constipation, blurred vision, dizziness and bloating. Less frequently occurring reactions include vivid dreams, impotence, tremor, confusion and nasal congestion. Many depressive symptoms including anorexia, fatigue, weakness, restlessness and lethargy have been reported as side effects of both Limbitrol and amitriptyline. Granulocytopenia, jaundice and hepatic dysfunction have been observed rarely.

The following list includes adverse reactions not reported with Limbitrol but requiring consideration because they have been reported with one or both components or closely related drugs. **Cardiovascular:** Hypotension, hypertension, tachycardia, palpitations, myocardial infarction, arrhythmias, heart block, stroke. **Psychiatric:** Euphoria, apprehension, poor concentration, delusions, hallucinations, hypomania and increased or decreased libido.

Neurologic: Incoordination, ataxia, numbness, tingling and paresthesias of the extremities, extrapyramidal symptoms, syncope, changes in EEG patterns.

Anticholinergic: Disturbance of accommodation, paralytic ileus, urinary retention, dilatation of urinary tract.

Allergic: Skin rash, urticaria, photosensitization, edema of face and tongue, pruritus.

Hematologic: Bone marrow depression including agranulocytosis, eosinophilia, purpura, thrombocytopenia.

Gastrointestinal: Nausea, epigastric distress, vomiting, anorexia, stomatitis, peculiar taste, diarrhea, black tongue.

Endocrine: Testicular swelling and gynecomastia in the male, breast enlargement, galactorrhea and minor menstrual irregularities in the female, elevation and lowering of blood sugar levels, and syndrome of inappropriate ADH (antidiuretic hormone) secretion.

Other: Headache, weight gain or loss, increased perspiration, urinary frequency, mydriasis, jaundice, alopecia, parotid swelling.

Overdosage: Immediately hospitalize patient suspected of having taken an overdose. Treatment is symptomatic and supportive. I.V. administration of 1 to 3 mg physostigmine salicylate has been reported to reverse the symptoms of amitriptyline poisoning. See complete product information for manifestation and treatment.

Dosage: Individualize according to symptom severity and patient response. Reduce to smallest effective dosage when satisfactory response is obtained. Larger portion of daily dose may be taken at bedtime. Single h.s. dose may suffice for some patients. Lower dosages are recommended for the elderly.

Limbitrol DS (double strength) Tablets, initial dosage of three or four tablets daily in divided doses, increased to six tablets or decreased to two tablets daily as required. Limbitrol Tablets, initial dosage of three or four tablets daily in divided doses, for patients who do not tolerate higher doses.

How Supplied: Double strength (DS) Tablets, white, film-coated, each containing 10 mg clordiazepoxide and 25 mg amitriptyline (as the hydrochloride salt), and Tablets, blue, film-coated, each containing 5 mg clordiazepoxide and 12.5 mg amitriptyline (as the hydrochloride salt)—bottles of 100 and 500; Tel-E-Dose[®] packages of 100; Prescription Paks of 50.



ROCHE PRODUCTS INC.
Manati, Puerto Rico 00701

In moderate depression and anxiety

CLEARING: BOTH DEPRESSION AND ANXIETY...



Now

IN PLACE OF
LIMBITROL 5-12.5 WRITE:

Limbitrol[®]

Each tablet contains 5 mg chlordiazepoxide and 12.5 mg amitriptyline (as the hydrochloride salt)



IN PLACE OF
LIMBITROL 10-25 WRITE:

Limbitrol DS[™]

Each tablet contains 10 mg chlordiazepoxide and 25 mg amitriptyline (as the hydrochloride salt)



Easier to remember...easier to prescribe

Copyright © 1985 by Roche Products Inc. All rights reserved.

Please see summary of product information on reverse side.